

PC

268.5

1155

no. 58

1978

National Cancer Institute

CARCINOGENESIS

Technical Report Series

No. 58

1978

**BIOASSAY OF
THIO-TEPA
FOR POSSIBLE CARCINOGENICITY**

CAS No. 52-24-4

NCI-CG-TR-58

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Institutes of Health



1864
1865
1866

2-10 7/29/78

BIOASSAY OF
THIO-TEPA
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program
Division of Cancer Cause and Prevention
National Cancer Institute
National Institutes of Health
Bethesda, Maryland 20014

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Institutes of Health

DHEW Publication No. (NIH) 78-1308

BIOASSAY OF
THIO-TEPA
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program
Division of Cancer Cause and Prevention
National Cancer Institute
National Institutes of Health

FOREWORD: This report presents the results of the bioassay of thio-TEPA conducted for the Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute (NCI), National Institutes of Health, Bethesda, Maryland. This is one of a series of experiments designed to determine whether selected chemicals have the capacity to produce cancer in animals. Negative results, in which the test animals do not have a greater incidence of cancer than control animals, do not necessarily mean that the test chemical is not a carcinogen, inasmuch as the experiments are conducted under a limited set of circumstances. Positive results demonstrate that the test chemical is carcinogenic for animals under the conditions of the test and indicate that exposure to the chemical is a potential risk to man. The actual determination of the risk to man from animal carcinogens requires a wider analysis.

CONTRIBUTORS: This bioassay of thio-TEPA was conducted by Southern Research Institute, Birmingham, Alabama, initially under direct contract to NCI and currently under a subcontract to Tracor Jitco, Inc., prime contractor for the NCI Carcinogenesis Testing Program.

The experimental design and doses were determined by Drs. D. P. Griswold¹, J. D. Prejean¹, E. K. Weisburger², and J. H. Weisburger^{2,3}. Ms. J. Belzer¹ and Mr. I. Brown¹ were responsible for the care of the laboratory animals and administration of the test chemical. Data management and retrieval were performed by Ms. C. A. Dominick¹. Histopathologic examinations were performed by Drs. S. D. Kosanke¹ and J. C. Peckham¹, and the diagnoses included in this report represent their interpretation.

Animal pathology tables and survival tables were compiled at EG&G Mason Research Institute⁴. The statistical analyses were performed by Dr. J. R. Joiner⁵ and Ms. P. L. Yong⁵, using methods selected for the bioassay program by Dr. J. J. Gart⁶. Chemicals used in this bioassay were analyzed under the direction of Dr. E. Murrill⁷, and the analytical results were reviewed by Dr. S. S. Olin⁵. The chemical structure was supplied by NCI.

This report was prepared at Tracor Jitco⁵ under the direction of NCI. Those responsible for the report at Tracor Jitco were Dr. L. A. Campbell, Director of the Bioassay Program; Dr. S. S. Olin, Deputy Director for Science; Dr. J. F. Robens, toxicologist; Dr. R. L. Schueler, pathologist; Dr. G. L. Miller, Ms. L. A. Waitz, and Mr. W. D. Reichardt, bioscience writers; and Dr. E. W. Gunberg, technical editor, assisted by Ms. Y. E. Presley.

The following other scientists at NCI were responsible for evaluating the bioassay experiment, interpreting the results, and reporting the findings: Dr. Kenneth C. Chu, Dr. Cipriano Cueto, Jr., Dr. J. Fielding Douglas, Dr. Dawn G. Goodman, Dr. Richard A. Griesemer, Dr. Harry A. Milman, Dr. Thomas W. Orme, Dr. Robert A. Squire⁸, Dr. Jerrold M. Ward, and Dr. Carrie E. Whitmire.

¹Southern Research Institute, 2000 Ninth Avenue South, Birmingham, Alabama.

²Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.

³Now with the Naylor Dana Institute for Disease Prevention, American Health Foundation, Hammond House Road, Valhalla, New York.

⁴EG&G Mason Research Institute, 1530 East Jefferson Street, Rockville, Maryland.

⁵Tracor Jitco, Inc., 1776 East Jefferson Street, Rockville, Maryland.

⁶Mathematical Statistics and Applied Mathematics Section,
Biometry Branch, Field Studies and Statistics, Division of
Cancer Cause and Prevention, National Cancer Institute, National
Institutes of Health, Bethesda, Maryland.

⁷Midwest Research Institute, 425 Volker Boulevard, Kansas City,
Missouri.

⁸Now with the Division of Comparative Medicine, Johns Hopkins
University, School of Medicine, Traylor Building, Baltimore,
Maryland.

SUMMARY

A bioassay of thio-TEPA for possible carcinogenicity was conducted by administering the test chemical by intraperitoneal injection to Sprague-Dawley rats and B6C3F1 mice.

Groups of 31-39 rats of each sex were administered thio-TEPA in phosphate-buffered saline at one of three doses, either 0.7, 1.4, or 2.8 mg/kg body weight, three times per week for a maximum of 52 weeks, then observed for additional periods of time. The maximum time on study (administration of chemical and observation) was 86 weeks. The groups at the low dose were started 69 weeks after those at the mid and high doses, because of high mortalities observed in the groups at the higher doses. Matched controls consisted of groups of 10 untreated rats and 10 vehicle-control rats of each sex. Pooled-control groups also were used. Surviving control rats were killed at 82-87 weeks; surviving dosed rats were killed at 81 or 82 weeks.

Groups of 35 mice of each sex were administered thio-TEPA at one of two doses, either 1.15 or 2.3 mg/kg body weight, three times per week for a maximum of 52 weeks, then observed for a maximum additional period of 34 weeks. Matched controls consisted of groups of 15 untreated mice and 15 vehicle-control mice of each sex. Pooled controls also were used. Surviving control and dosed mice were killed at 86 or 87 weeks.

Thio-TEPA was toxic to both rats and mice, causing decreased mean body weight gains and early deaths in the mid- and high-dose rats and in the high-dose mice. Because of the early deaths, statistical analyses were based only on time-adjusted incidences of tumors. Since all high-dose male and female rats had died by 21 weeks, microscopic evaluation of tissues was performed only on the low- and mid-dose animals.

In rats, the incidence of combined neoplasms of the hematopoietic system (lymphoma, lymphocytic leukemia, or granulocytic leukemia) was significant in the males in both the low-dose ($P = 0.020$) and mid-dose ($P = 0.001$) groups, using pooled controls (pooled

controls 0/29, low-dose 6/34; pooled controls 0/30, mid-dose 6/16).

Squamous-cell carcinoma of the skin or ear canal occurred at a significant incidence in the male rats in both the low-dose ($P = 0.009$) and mid-dose ($P = 0.023$) groups, using pooled controls (pooled controls 0/29, low-dose 7/33; pooled controls 0/30, mid-dose 3/13) and in the mid-dose females ($P < 0.001$), using pooled controls (pooled controls 0/28, mid-dose 8/21); in addition, two low-dose females had such tumors, with none occurring in the corresponding low-dose controls.

The incidence of adenocarcinoma of the uterus was significant in the mid-dose female rats ($P = 0.001$), using pooled controls (pooled controls 0/28, mid-dose 7/21); in addition, two low-dose females had adenocarcinoma of the uterus, with no such tumor occurring in the corresponding low-dose controls.

In rats, neuroepitheliomas (neuroblastomas) or nasal carcinomas occurred in three low-dose males, two low-dose females, and two mid-dose females. Although these are not statistically significant incidences, these tumors did not occur among control animals and no such tumors have occurred in 380 Sprague-Dawley control rats of each sex in other bioassays at the same laboratory. Thus, they may be associated with administration of the chemical.

In the high-dose groups of both male and female mice, but not in the low-dose groups, the incidences of lymphoma or lymphocytic leukemia were significantly higher ($P < 0.001$) for each sex than those of either the vehicle or pooled controls (males: vehicle controls 1/8, pooled controls 1/18, low-dose 2/24, high-dose 26/28; females: vehicle controls 0/14, pooled controls 0/29, low-dose 5/26, high-dose 32/32).

In the low-dose male mice squamous-cell carcinoma was found in the skin of seven animals, in the preputial glands of six

animals, and in the ear canal of two animals. A carcinoma of the preputial gland was also found in a high-dose male. When the incidences of the tumors at the different sites were combined, the incidence in the low-dose group was statistically significant using either the vehicle ($P = 0.004$) or the pooled ($P < 0.001$) controls (vehicle controls 0/8, pooled controls 0/18, low-dose 14/24, high-dose 1/2).

It is concluded that under the conditions of this bioassay, thio-TEPA was carcinogenic in both Sprague-Dawley rats and B6C3F1 mice. In the rats, the chemical induced squamous-cell carcinoma of the skin or ear canal in both males and females, and hematopoietic neoplasms in the males; in the mice, it induced lymphoma or lymphocytic leukemia in both sexes and squamous-cell carcinoma in the skin and associated glands of males.

TABLE OF CONTENTS

	<u>Page</u>
I. Introduction.....	1
II. Materials and Methods.....	3
A. Chemical.....	3
B. Dosage Preparation.....	4
C. Animals.....	4
D. Animal Maintenance.....	5
E. Subchronic Studies.....	8
F. Designs of Chronic Studies.....	9
G. Clinical and Pathologic Examinations.....	14
H. Data Recording and Statistical Analyses.....	15
III. Results - Rats.....	21
A. Body Weights and Clinical Signs (Rats).....	21
B. Survival (Rats).....	21
C. Pathology (Rats).....	26
D. Statistical Analyses of Results (Rats).....	33
IV. Results - Mice.....	37
A. Body Weights and Clinical Signs (Mice).....	37
B. Survival (Mice).....	37
C. Pathology (Mice).....	40
D. Statistical Analyses of Results (Mice).....	43
V. Discussion.....	45
VI. Bibliography.....	49

APPENDIXES

Appendix A	Summary of the Incidence of Neoplasms in Rats Given Intraperitoneal Injections of Thio-TEPA.....	51
Table A1	Summary of the Incidence of Neoplasms in Male Rats Given Intraperitoneal Injections of Thio-TEPA (Control Groups)...	53

	<u>Page</u>
Table A2	Summary of the Incidence of Neoplasms in Male Rats Given Intraperitoneal Injections of Thio-TEPA (Treated Groups)... 56
Table A3	Summary of the Incidence of Neoplasms in Female Rats Given Intraperitoneal Injections of Thio-TEPA (Control Groups)... 60
Table A4	Summary of the Incidence of Neoplasms in Female Rats Given Intraperitoneal Injections of Thio-TEPA (Treated Groups)... 63
Appendix B	Summary of the Incidence of Neoplasms in Mice Given Intraperitoneal Injections of Thio-TEPA..... 67
Table B1	Summary of the Incidence of Neoplasms in Male Mice Given Intraperitoneal Injections of Thio-TEPA..... 69
Table B2	Summary of the Incidence of Neoplasms in Female Mice Given Intraperitoneal Injections of Thio-TEPA..... 72
Appendix C	Summary of the Incidence of Nonneoplastic Lesions in Rats Given Intraperitoneal Injections of Thio-TEPA..... 75
Table C1	Summary of the Incidence of Nonneoplastic Lesions in Male Rats Given Intraperitoneal Injections of Thio-TEPA (Control Groups)... 77
Table C2	Summary of the Incidence of Nonneoplastic Lesions in Male Rats Given Intraperitoneal Injections of Thio-TEPA (Treated Groups)... 80
Table C3	Summary of the Incidence of Nonneoplastic Lesions of Female Rats Given Intraperitoneal Injections of Thio-TEPA (Control Groups)... 84
Table C4	Summary of the Incidence of Nonneoplastic Lesions of Female Rats Given Intraperitoneal Injections of Thio-TEPA (Treated Groups)... 87

	<u>Page</u>
Appendix D	Summary of the Incidence of Nonneoplastic Lesions in Mice Given Intraperitoneal Injections of Thio-TEPA..... 91
Table D1	Summary of the Incidence of Nonneoplastic Lesions in Male Mice Given Intraperitoneal Injections of Thio-TEPA..... 93
Table D2	Summary of the Incidence of Nonneoplastic Lesions in Female Mice Given Intraperitoneal Injections of Thio-TEPA..... 96
Appendix E	Analyses of the Incidence of Primary Tumors in Rats Given Intraperitoneal Injections of Thio-TEPA..... 99
Table E1	Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA....101
Table E2	Analyses of the Incidence of Primary Tumors in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA....107
Table E3	Analyses of the Incidence of Primary Tumors in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA....110
Table E4	Analyses of the Incidence of Primary Tumors in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA....116
Table E5	Time-adjusted Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA.....;.....121
Table E6	Time-adjusted Analyses of the Incidence of Primary Tumors in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA.....127

	<u>Page</u>
Table E7	Time-adjusted Analyses of the Incidence of Primary Tumors in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA.....130
Table E8	Time-adjusted Analyses of the Incidence of Primary Tumors in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA.....136
Appendix F	Analyses of the Incidence of Primary Tumors in Mice Given Intraperitoneal Injections of Thio-TEPA.....143
Table F1	Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA.....145
Table F2	Analyses of the Incidence of Primary Tumors in Female Mice Given Intraperitoneal Injections of Thio-TEPA.....154
Table F3	Time-adjusted Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA....158
Table F4	Time-adjusted Analyses of the Incidence of Primary Tumors in Female Mice Given Intraperitoneal Injections of Thio-TEPA....164

TABLES

Table 1	Design of Chronic Studies of Thio-TEPA in Rats..... 10
Table 2	Design of Chronic Studies of Thio-TEPA in Mice..... 12

FIGURES

Figure 1	Growth Curves for Male Rats Treated with Thio-TEPA..... 22
Figure 2	Growth Curves for Female Rats Treated with Thio-TEPA..... 23

	<u>Page</u>
Figure 3	Survival Curves for Male Rats Treated with Thio-TEPA..... 24
Figure 4	Survival Curves for Female Rats Treated with Thio-TEPA..... 25
Figure 5	Growth Curves for Mice Treated with Thio-TEPA..... 38
Figure 6	Survival Curves for Mice Treated with Thio-TEPA..... 39




I. INTRODUCTION

Thio-TEPA (CAS 52-24-4; NSC 6396; NCI C01649) is an ethyleneimine alkylating agent that was introduced in 1953 for clinical use in cancer chemotherapy. Following oxidative desulfurization of thio-TEPA, the three ethylenimine groups that the chemical contains become activated to ethylenimonium ions (Montgomery and Struck, 1973; Lederle Laboratories, 1977). The presence of several such groups raises the possibility that the drug will bind to more than one site on the DNA molecule and will have increased cytotoxic effects (Calabresi and Parks, 1975).

At one time thio-TEPA was an important therapeutic drug in the management of ovarian carcinoma (Calabresi and Welch, 1962). It has been used effectively in the treatment of Hodgkins disease, bronchogenic carcinoma, bladder cancer, retinoblastoma, and breast cancer and for the control of pleural, pericardial, and peritoneal neoplastic effusions (Wheeler, 1973; Carter and Slavik, 1974; Calabresi and Parks, 1975). In the pulmonary tumor response test in strain A mice (Shimkin et al., 1966), thio-TEPA showed weak activity. On the basis of these results, the drug was retested in a lifetime carcinogen bioassay. This drug is one of a series of anticancer agents that were selected for testing because they may be administered chronically in humans.

II. MATERIALS AND METHODS

A. Chemical



Thio-TEPA

Thio-TEPA is the common name for tris(1-aziridinyl)phosphine sulfide, manufactured by Lederle Laboratories, Pearl River, New York. Lederle Lot No. D9878, which conformed to USP specifications, was used during the chronic study. The purity of this lot was $98.0 \pm 1.0\%$ as determined by titration of the aziridine groups with tetrabutylammonium iodide and perchloric acid. Elemental analysis for sulfur was slightly high ($18.2 \pm 0.6\%$ vs. 17.0% theoretical), but may have been affected by phosphorus; analysis of carbon, hydrogen, nitrogen, and phosphorus verified the chemical composition of this product. The melting point was $52.0-52.5^{\circ}\text{C}$ and was comparable to that of 51.5°C reported elsewhere (U.S. Patent, 1954). Thin-layer

chromatography showed one trace impurity which remained at the origin. Vapor phase chromatography gave one homogeneous peak. The infrared and nuclear magnetic resonance spectra were consistent with the spectra given in the literature for this compound.

B. Dosage Preparation

Solutions of thio-TEPA in phosphate-buffered saline were prepared on the day that they were used by blending for 20 seconds in a 10-ml Potter-Elvehjem tissue grinder with a Teflon pestle; unused solutions of the drug were discarded. Concentrations of thio-TEPA ranged from 0.028% to 0.112% for the chronic studies using rats and from 0.012% to 0.023% for the chronic studies using mice. Volumes of solutions used for injection were 0.25 ml/100 g body weight for the rats and 1.0 ml/100 g body weight for the mice.

C. Animals

For the subchronic studies, male Sprague-Dawley rats and male Swiss mice were obtained from Charles River Breeding Laboratories, Inc., Wilmington, Massachusetts. On arrival at the laboratory, they were 30 days of age. The rats were quarantined for 5 days and the mice for 20 days prior to being placed on study.

In the chronic studies, Sprague-Dawley rats and B6C3F1 mice of each sex were obtained from Charles River Breeding Laboratories under a contract with the Division of Cancer Treatment, National Cancer Institute. Male rats were received at 29 days of age, female rats at 36 days of age, and male and female mice at 32 days of age. All animals were quarantined for periods of 6 days for rats and 10 days for mice. After these periods, animals with no visible signs of disease were assigned to control or dosed groups and earmarked for individual identification. Additional groups of male and female Sprague-Dawley rats for the chronic study were received at 30 days of age from Charles River Breeding Laboratories and quarantined for 28 days weeks prior to being placed on study.

D. Animal Maintenance

All animals were housed in temperature- and humidity-controlled rooms. The temperature range was 20-24°C, and the relative humidity was maintained at 40-60%. The air was changed 15 times per hour and passed through both intake and exhaust fiberglass roughing filters. In addition to natural light, illumination was provided by fluorescent light for 9 hours per day. Wayne® Lab Blox animal meal (Allied Mills, Inc., Chicago, Ill.) and water were supplied daily and were available ad libitum.

Rats were housed five per cage and mice seven per cage in solid-bottom stainless steel cages (Hahn Roofing and Sheet Metal Co., Birmingham, Ala.). Rat cages were provided with Iso-Dri[®] hardwood chip bedding (Carworth, Edison, N.J.); mouse cages were provided with Sterolit[®] clay bedding (Englehard Mineral and Chemical Co., New York, N.Y.). For initial studies with rats, cage tops were covered with disposable filter bonnets beginning at week 24; for later studies with rats, cage tops were covered with filter bonnets for the entire test. Filter bonnets were not used for the mouse cages. Bedding was replaced once per week; cages, water bottles, and feeders were sanitized at 82°C once per week; and racks were cleaned once per week.

The rats and mice were housed in separate rooms. Control animals were housed with respective dosed animals. Animals administered thio-TEPA were maintained in the same rooms as animals of the same species being administered the following chemicals:

RATS

Gavage Studies

cholesterol (p-(bis(2-chloroethyl)amino)phenyl)acetate
(phenesterin) (CAS 3546-10-9)
estradiol bis((p-(bis(2-chloroethyl)amino)phenyl)acetate)
(estradiol mustard) (CAS 22966-79-6)

Intraperitoneal Injection Studies

4'-(9-acridinylamino)methansulfon-m-aniside monohydrochloride
(MAAM) (NSC 141549)

acronycine (CAS 7008-42-6)
5-azacytidine (CAS 320-67-2)
beta-2'-deoxy-6-thioguanosine monohydrate (beta-TGdR)
(CAS 789-61-7)
1,4-butanediol dimethanesulfonate (busulfan) (CAS 55-98-1)
emetine dihydrochloride tetrahydrate (CAS 316-42-7)
3,3'-iminobis-1-propanol dimethanesulfonate (ester)
hydrochloride [IPD] (CAS 3458-22-8)
(+)-4,4'-(1-methyl-1,2-ethanediyl)bis-2,6-piperazinedione
(ICRF-159) (CAS 21416-87-5)
N,3-bis(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-
amine-2-oxide (isophosphamide) (CAS 3778-73-2)
N-(2-chloroethyl)-N-(1-methyl-2-phenoxyethyl)benzylamine
hydrochloride (phenoxybenzamine) (CAS 63-92-3)
N-(1-methylethyl)-4-((2-methylhydrazino)methyl)benzamide
monohydrochloride (procarbazine) (CAS 366-70-1)
2,4,6-tris(dimethylamino)-s-triazine (CAS 645-05-6)
adriamycin (CAS 23214-92-8)

MICE

Feed Studies

4-acetyl-N-((cyclohexylamino)carbonyl)benzenesulfonamide
(acetohexamide) (CAS 968-81-0)
anthranilic acid (CAS 118-92-3)
1-butyl-3-(p-tolylsulfonyl)urea (tolbutamide) (CAS 64-77-7)
4-chloro-N-((propylamino)carbonyl)benzenesulfonamide
(chlorpropamide) (CAS 94-20-2)
5-(4-chlorophenyl)-6-ethyl-2,4-pyrimidinediamine
(pyrimethamine) (CAS 58-14-0)
2,6-diamino-3-(phenylazo)pyridine hydrochloride
(phenazopyridine hydrochloride) (CAS 136-40-3)
L-tryptophan (CAS 73-22-3)
N-9H-fluoren-2-ylacetamide (CAS 53-96-3)
N-(p-toluenesulfonyl)-N'-hexamethyleniminourea
(tolazamide) (CAS 1156-19-0)
1-phenethylbiguanide hydrochloride (phenformin) (CAS 114-86-3)
pyrazinecarboxamide (pyrazinamide) (CAS 98-96-4)
4,4'-sulfonyldianiline (dapsone) (CAS 80-08-0)
4,4'-thiodianiline (CAS 139-65-1)
ethionamide (CAS 536-33-4)

Gavage Studies

cholesterol (p-(bis(2-chloroethyl)amino)phenyl)acetate
(phenesterin) (CAS 3546-10-9)

estradiol bis((p-(bis(2-chloroethyl)amino)phenyl)acetate)
(estradiol mustard) (CAS 22966-79-6)

Intraperitoneal Injection Studies

4'-(9-acridinylamino)methansulfon-m-aniside monohydrochloride
(MAAM) (NSC 141549)
acronycine (CAS 7008-42-6)
5-azacytidine (CAS 320-67-2)
beta-2'-deoxy-6-thioguanosine monohydrate (beta-TGdR)
(CAS 789-61-7)
1,4-butanediol dimethanesulfonate (busulfan) (CAS 55-98-1)
emetine dihydrochloride tetrahydrate (CAS 316-42-7)
3,3'-iminobis-1-propanol dimethanesulfonate (ester)
hydrochloride [IPD] (CAS 3458-22-8)
(+)-4,4'-(1-methyl-1,2-ethanediyl)bis-2,6-piperazinedione
(ICRF-159) (CAS 21416-87-5)
N,3-bis(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-
amine-2-oxide (isophosphamide) (CAS 3778-73-2)
N-(2-chloroethyl)-N-(1-methyl-2-phenoxyethyl)benzylamine
hydrochloride (phenoxybenzamine) (CAS 63-92-3)
N-(1-methylethyl)-4-((2-methylhydrazino)methyl)benzamide
monohydrochloride (procarbazine) (CAS 366-70-1)

E. Subchronic Studies

Subchronic studies were conducted using male Sprague-Dawley rats and male Swiss mice to estimate the maximum tolerated doses of thio-TEPA, on the basis of which two concentrations (hereinafter referred to as "low doses" and "high doses") were determined for administration in the chronic studies. Rats were administered the drug at doses of 0.14, 0.35, 0.7, 1.4, or 2.8 mg/kg, and mice at doses of 0.23, 0.58, 1.15, 2.3, or 4.6 mg/kg. Dosed animals were injected intraperitoneally with thio-TEPA three times per week for 45 days and then observed for an additional 45 days. Five animals of each species were administered the chemical at

each dose, 10 animals of each species were maintained as vehicle controls, and 10 animals of each species were maintained as untreated controls.

There were no deaths in the rats at any dose tested. Mean body weight gains at 45 days in animals at 0.14, 0.35, or 0.7 mg/kg were unaffected in comparison with those of the vehicle controls; at 1.4 mg/kg the mean body weights were 79% of those of the controls, and at 2.8 mg/kg they were 70% of those of the controls. After 90 days, mean body weight gains were comparable to those of the controls in all groups except those at 1.4 mg/kg and 2.8 mg/kg, where they were approximately 85% of those of the controls. The vehicle controls gained slightly less weight than the untreated controls. The low and high doses for the chronic studies using rats were set at 1.4 and 2.8 mg/kg.

In mice, 3/5 animals died at 4.6 mg/kg and 1/5 died at 0.58 mg/kg. At 45 days, mean body weight gains in dosed animals were not greatly affected, except for slight losses in the group at 4.6 mg/kg. The low and high doses for the chronic studies using mice were set at 1.15 and 2.3 mg/kg.

F. Designs of the Chronic Studies

The designs of the chronic studies are shown in tables 1 and 2.

Table 1. Design of Chronic Studies of Thio-TEPA in Rats

Sex and Test Group	Initial No. of Animals ^a	Thio-TEPA Dose ^b (mg/kg)	Time on Study	
			Dosed (weeks)	Observed (weeks)
<u>Male</u>				
Low-Dose Untreated-Control ^c	10	0		83
Low-Dose Vehicle-Control ^c	10	0 ^d	52	30
Low-Dose ^c	39	0.7	52	30
Mid- and High-Dose Untreated-Control	10	0		87
Mid- and High-Dose Vehicle-Control	10	0 ^d	34 ^e	52
Mid-Dose	35	1.4	34 ^e	44 ^f
High-Dose	35	2.8	19 ^g	
<u>Female</u>				
Low-Dose Untreated-Control ^c	10	0		82
Low-Dose Vehicle-Control ^c	10	0 ^d	52	30
Low-Dose ^c	31	0.7	52	30
Mid- and High-Dose Untreated-Control	10	0		87
Mid- and High-Dose Vehicle-Control	10	0 ^d	34 ^e	53
Mid-Dose	35	1.4	34 ^e	47
High-Dose	35	2.8	21 ^g	

^aAges of rats when placed on study: mid- and high-dose males, 35 days; mid- and high-dose females, 42 days; low-dose males and females, 58 days.

^bThio-TEPA was administered intraperitoneally in phosphate-buffered saline three times per week at a volume of 0.25 ml/100 g body weight; doses were based on individual weights.

Table 1. Design of Chronic Studies of Thio-TEPA in Rats

(continued)

^cBecause of deaths in the high- and mid-dose groups, new dosed and control groups were started 69 weeks after the start of the original study.

^dVehicle controls received phosphate-buffered saline at the same volume as the dosed animals.

^eAdministration of the chemical to mid-dose males and females and to mid- and high-dose vehicle controls terminated at week 34, due to toxicity in the dosed groups.

^fObservation of mid-dose males terminated at week 78, due to death of all animals.

^gAdministration of the chemical to high-dose males and females terminated at times indicated, due to death of all animals.

Table 2. Design of Chronic Studies of Thio-TEPA in Mice

Sex and Test Group	Initial No. of Animals ^a	Thio-TEPA Dose ^b (mg/kg)	Time on Study	
			Dosed (weeks)	Observed (weeks)
<u>Male</u>				
Untreated-Control	15	0		87
Vehicle-Control	15	0 ^c	52	35
Low-Dose	35	1.15	52	34
High-Dose	35	2.3	52	4 ^d
<u>Female</u>				
Untreated-Control	15	0		87
Vehicle-Control	15	0 ^c	52	35
Low-Dose	35	1.15	52	34
High-Dose	35	2.3	43 ^e	

^aAll animals were 42 days of age when placed on study.

^bThio-TEPA was administered intraperitoneally in phosphate-buffered saline three times per week at a volume of 1.0 ml/100 g body weight based on the mean weight of the animals in each cage.

^cVehicle controls received only phosphate-buffered saline solution, at the same volume as dosed mice.

^dObservation of high-dose males terminated at week 56, due to death of all animals.

^eAdministration of the chemical to high-dose females terminated at week 43, due to death of all animals.

Since the numbers of rats and mice in the control groups were small, pooled-control groups also were used for statistical comparisons. Pooled-control groups for the low-dose rats consisted of vehicle-control groups of 10 animals of each sex from the bioassay of thio-TEPA, combined with corresponding vehicle-control groups of 10 animals of each sex from similar bioassays of beta-2'-deoxy-6-thioguanosine monohydrate (β -TGDR) and 3,3'-iminobis-1-propanol dimethanesulfonate (ester) [IPD]; pooled-control groups for the mid-dose rats consisted of vehicle-control groups of 10 animals of each sex from the bioassay of thio-TEPA, combined with corresponding vehicle-control groups of 10 animals of each sex from similar bioassays of procarbazine and IPD. The total number of animals was 30 in both the low-dose pooled-control group and the mid-dose pooled-control group. For the mice, pooled-control groups consisted of vehicle-control groups of 15 animals of each sex from the bioassay of thio-TEPA, combined with corresponding vehicle-control groups of 15 animals of each sex from a similar bioassay of IPD, to give a total of 30 animals per group. The bioassays of the chemicals other than thio-TEPA were also conducted at Southern Research Institute and were started no more than 3 months apart from those for thio-TEPA. The vehicle-control groups of rats and mice used in the pooled-control groups were of the same strain, obtained from the same supplier, and examined by the same pathologists as in the

respective dosed groups; further, the same vehicle was used for the different vehicle-control groups.

G. Clinical and Pathologic Examinations

All animals were observed twice daily for signs of toxicity, and animals that were moribund were killed and necropsied, except for those dying prior to day 100, due, presumably, to toxicity of the test chemical. Rats (mid- and high-dose) and mice were weighed individually each week for 8 weeks and every 2 weeks thereafter; low-dose rats were weighed once every 2 weeks for the period of administration of the chemical and once per month thereafter. Palpation for masses was carried out at each weighing.

The pathologic evaluation consisted of gross and microscopic examination of major tissues, major organs, and all gross lesions from killed animals and from animals found dead. The following tissues were examined microscopically: skin, muscle, lungs and bronchi, trachea, bone and bone marrow, spleen, lymph nodes, thymus, heart, salivary gland, liver, gallbladder and bile duct (mice), pancreas, esophagus, stomach, small intestine, large intestine, kidney, urinary bladder, pituitary, adrenal, thyroid, parathyroid, mammary gland, prostate or uterus, testis or ovary, brain, and sensory organs. Peripheral blood smears were prepared from each animal whenever possible. Occasionally, additional

tissues were also examined microscopically. The different tissues were preserved in 10% buffered formalin, embedded in paraffin, sectioned, and stained with hematoxylin and eosin. Special staining techniques were utilized when indicated for more definitive diagnosis.

A few tissues from some animals were not examined, particularly from those animals that died early. Also, some animals were missing, cannibalized, or judged to be in such an advanced state of autolysis as to preclude histopathologic evaluation. Thus, the number of animals from which particular organs or tissues were examined microscopically varies, and does not necessarily represent the number of animals that were placed on study in each group.

H. Data Recording and Statistical Analyses

Pertinent data on this experiment have been recorded in an automatic data processing system, the Carcinogenesis Bioassay Data System (Linhart et al., 1974). The data elements include descriptive information on the chemicals, animals, experimental design, clinical observations, survival, body weight, and individual pathologic results, as recommended by the International Union Against Cancer (Berenblum, 1969). Data

tables were generated for verification of data transcription and for statistical review.

These data were analyzed using the statistical techniques described in this section. Those analyses of the experimental results that bear on the possibility of carcinogenicity are discussed in the statistical narrative sections.

Probabilities of survival were estimated by the product-limit procedure of Kaplan and Meier (1958) and are presented in this report in the form of graphs. Animals were statistically censored as of the time that they died of other than natural causes or were found to be missing; animals dying from natural causes were not statistically censored. Statistical analyses for a possible dose-related effect on survival used the method of Cox (1972) for testing two groups for equality and Tarone's (1975) extensions of Cox's methods for testing for a dose-related trend. One-tailed P values have been reported for all tests except the departure from linearity test, which is only reported when its two-tailed P value is less than 0.05.

The incidence of neoplastic or nonneoplastic lesions has been given as the ratio of the number of animals bearing such lesions at a specific anatomic site (numerator) to the number of animals in which that site is examined (denominator). In most instances,

the denominators included only those animals for which that site was examined histologically. However, when macroscopic examination was required to detect lesions prior to histologic sampling (e.g., skin or mammary tumors), or when lesions could have appeared at multiple sites (e.g., lymphomas), the denominators consist of the numbers of animals necropsied.

The purpose of the statistical analyses of tumor incidence is to determine whether animals receiving the test chemical developed a significantly higher proportion of tumors than did the control animals. As a part of these analyses, the one-tailed Fisher exact test (Cox, 1970) was used to compare the tumor incidence of a control group with that of a group of dosed animals at each dose level. When results for a number of dosed groups (k) are compared simultaneously with those for a control group, a correction to ensure an overall significance level of 0.05 may be made. The Bonferroni inequality (Miller, 1966) requires that the P value for any comparison be less than or equal to $0.05/k$. In cases where this correction was used, it is discussed in the narrative section. It is not, however, presented in the tables, where the Fisher exact P values are shown.

The Cochran-Armitage test for linear trend in proportions, with continuity correction (Armitage, 1971), was also used. Under the assumption of a linear trend, this test determines if the slope

of the dose-response curve is different from zero at the one-tailed 0.05 level of significance. Unless otherwise noted, the direction of the significant trend is a positive dose relationship. This method also provides a two-tailed test of departure from linear trend.

A time-adjusted analysis was applied when numerous early deaths resulted from causes that were not associated with the formation of tumors. In this analysis, deaths that occurred before the first tumor was observed were excluded by basing the statistical tests on animals that survived at least 52 weeks, unless a tumor was found at the anatomic site of interest before week 52. When such an early tumor was found, comparisons were based exclusively on animals that survived at least as long as the animal in which the first tumor was found. Once this reduced set of data was obtained, the standard procedures for analyses of the incidence of tumors (Fisher exact tests, Cochran-Armitage tests, etc.) were followed.

When appropriate, life-table methods were used to analyze the incidence of tumors. Curves of the proportions surviving without an observed tumor were computed as in Saffiotti et al. (1972). The week during which an animal died naturally or was sacrificed was entered as the time point of tumor observation. Cox's methods of comparing these curves were used for two groups;

Tarone's extension to testing for linear trend was used for three groups. The statistical tests for the incidence of tumors which used life-table methods were one-tailed and, unless otherwise noted, in the direction of a positive dose relationship. Significant departures from linearity ($P < 0.05$, two-tailed test) were also noted.

The approximate 95 percent confidence interval for the relative risk of each dosed group compared with its control was calculated from the exact interval on the odds ratio (Gart, 1971). The relative risk is defined as p_t/p_c where p_t is the true binomial probability of the incidence of a specific type of tumor in a dosed group of animals and p_c is the true probability of the spontaneous incidence of the same type of tumor in a control group. The hypothesis of equality between the true proportion of a specific tumor in a dosed group and the proportion in a control group corresponds to a relative risk of unity. Values in excess of unity represent the condition of a larger proportion in the dosed group than in the control.

The lower and upper limits of the confidence interval of the relative risk have been included in the tables of statistical analyses. The interpretation of the limits is that in approximately 95% of a large number of identical experiments, the true ratio of the risk in a dosed group of animals to that in a

control group would be within the interval calculated from the experiment. When the lower limit of the confidence interval is greater than one, it can be inferred that a statistically significant result ($P < 0.025$ one-tailed test when the control incidence is not zero, $P < 0.050$ when the control incidence is zero) has occurred. When the lower limit is less than unity, but the upper limit is greater than unity, the lower limit indicates the absence of a significant result while the upper limit indicates that there is a theoretical possibility of the induction of tumors by the test chemical, which could not be detected under the conditions of this test.

III. RESULTS - RATS

A. Body Weights and Clinical Signs (Rats)

Mean body weights of all dosed male rats, particularly those of the mid- and high-dose groups, were depressed throughout the study, when compared with either matched or vehicle controls; those of the dosed females were less markedly depressed (figures 1 and 2). Fluctuations in the growth curve may be due to mortality; as the size of a group diminishes, the mean body weight may be subject to wide variation.

There was clinical evidence of respiratory disease in the high- and mid-dose groups of rats and their untreated and vehicle controls. To control the respiratory disease, these animals received oxytetracycline in drinking water at a dose of 0.6 mg/ml during weeks 24-30 and 0.3 mg/ml during weeks 30-35. No other signs of toxicity were reported for the rats.

B. Survival (Rats)

The Kaplan and Meier curves estimating the probabilities of survival for the male and female rats administered thio-TEPA at the doses of this bioassay, together with those of the controls, are shown in figures 3 and 4.

For each sex, the results of the Tarone test were significant




Figure 1. Growth Curves for Male Rats Treated with Thio-TEPA






Figure 2. Growth Curves for Female Rats Treated with Thio-TEPA







Figure 4. Survival Curves for Female Rats Treated with Thio-TEPA

($P < 0.001$), indicating that the administration of thio-TEPA decreased survival in a dose-related manner. At the high dose, all animals were dead at week 19 (males) or week 21 (females). Survival at the mid dose was also low, with all males dead by week 78, and with only 3/35 (8.6%) females surviving to termination, while 6/39 (15.4%) low-dose males and 13/31 (42%) low-dose females survived to the end of the study. Among the male vehicle controls, 8/10 (80%) of the mid- and high-dose controls and 10/10 (100%) of the low-dose vehicle controls survived to week 82; corresponding values for females were 7/10 (70%) and 9/10 (90%). Even using time-adjusted analyses, eliminating those rats dying before week 52 on study, in some dosed groups, insufficient numbers of animals were at risk for the development of late-appearing tumors.

C. Pathology (Rats)

Histopathologic findings on neoplasms in rats are summarized in Appendix A, tables A1-A4; findings on nonneoplastic lesions are summarized in Appendix C, tables C1-C4.

A variety of neoplasms were observed in the untreated controls, vehicle controls (phosphate-buffered saline), and dosed groups. Some types of neoplasms occurred only, or with a greater frequency, in rats of dosed groups as compared with controls.

Most of these lesions, however, are not uncommon in this strain of rat independent of the administration of any chemical. The incidence of several tumors was higher in the dosed groups. These tumors involved the skin, connective tissues, ear canal, nasal cavity and brain, hematopoietic tissues, lymphoreticular tissues, and uterus. The incidences of these tumors were as follows:

	Combined ^a Untreated Controls	Combined ^a Vehicle Controls	Low Dose	Mid Dose
<u>MALES</u>				
Number of rats necropsied	(19)	(20)	(37)	(30)
<u>Integument</u>				
Squamous-cell papilloma	0	1	1	0
Squamous-cell carcinoma	0	0	5	3
Trichoepithelioma	0	0	1	0
Basal-cell carcinoma	0	0	0	1
Sarcoma, NOS ^b	0	0	5	0
Fibroma	0	2	2	0
Fibrosarcoma	0	0	1	0

^aMid- and high-dose control group and low-dose group control were combined.

^bNot otherwise specified

	<u>Combined Untreated Controls</u>	<u>Combined Vehicle Controls</u>	<u>Low Dose</u>	<u>Mid Dose</u>
Number of rats necropsied	(19)	(20)	(37)	(30)
<u>Ear Canal</u>				
Squamous-cell papilloma	0	0	1	0
Squamous-cell carcinoma	0	0	3	1
<u>Nasal Cavity</u>				
Carcinoma, NOS	0	0	1	0
<u>Hematopoietic and Lymphoreticular Tissues</u>				
Malignant lymphoma, histiocytic type	0	0	1	0
Leukemia, lymphocytic	0	0	3	5
Leukemia, granulocytic	0	0	2	1
<u>Brain^c</u>	(18)	(20)	(36)	(28)
Sarcoma, NOS	0	0	0	1
Astrocytoma	0	0	1	0
Neuroepithelioma (Neuroblastoma)	0	0	2	0
<u>Other Sites</u>	(19)	(20)	(37)	(30)
Sarcoma, NOS	0	0	4	1
<u>FEMALES</u>				
Number of rats necropsied	(20)	(20)	(30)	(33)
<u>Integument</u>				
Squamous-cell papilloma	0	0	0	1
Squamous-cell carcinoma	0	0	0	3
Sarcoma, NOS	0	0	1	0
Fibroma	1	0	1	0
Fibrosarcoma	0	0	1	0

^cNumber of rats with tissue examined microscopically

	<u>Combined Untreated Controls</u>	<u>Combined Vehicle Controls</u>	<u>Low Dose</u>	<u>Mid Dose</u>
Number of rats necropsied	(20)	(20)	(30)	(33)
<u>Ear Canal</u>				
Squamous-cell papilloma	0	0	0	1
Squamous-cell carcinoma	0	0	2	5
<u>Nasal Cavity</u>				
Carcinoma, NOS	0	0	0	1
<u>Brain^C</u>	(20)	(19)	(30)	(32)
Neuroepithelioma (Neuroblastoma)	0	0	2	1
Sarcoma, NOS	0	0	0	1
<u>Uterus^C</u>	(20)	(20)	(30)	(32)
Adenocarcinoma, NOS	0	0	2	7
Squamous-cell carcinoma	0	0	1	0
Sarcoma, NOS	0	0	2	1
Stromal polyp	0	0	4	1
<u>Other Sites^C</u>	(20)	(20)	(30)	(33)
Sarcoma, NOS	0	0	0	1

^CNumber of rats with tissue examined microscopically

The skin tumors were differentiated epithelial tumors with the majority being keratinizing squamous-cell carcinomas located in various regions of the head and neck. The squamous-cell tumors varied from benign acanthotic, hyperkeratotic, and papillary growths (papillomas) to less differentiated carcinomas that invaded the dermis. These latter tumors contained nests of

squamous cells forming whorls of keratin and cords or small groups of darkly staining anaplastic epithelial cells. The predominant cells were stratified and polygonal-shaped, had ample eosinophilic cytoplasm, and had large vesicular nuclei with prominent nucleoli. Two tumors appeared to arise from the basal cells of hair follicles, sebaceous glands, or other adnexal structures. The basal-cell tumors were composed of closely packed, small polygonal cells with scanty cytoplasm and darkly staining purple or blue round to oval nuclei. The cells had a tendency to form palisades. Various degrees of keratinization and hair-follicle formation were present. These tumors were classified as a trichoepithelioma, and a basal-cell carcinoma.

Several subcutaneous connective tissue tumors were observed in both dosed and untreated groups. These were spindle-cell tumors that varied from well-differentiated fibromas with extensive collagenous formation to less differentiated fibrosarcomas with less collagen and poorly differentiated sarcomas (sarcomas, NOS) with little or no collagen and irregular cystic or vascular spaces. The sarcomas occurred frequently around the head and neck as well as other subcutaneous and visceral sites.

Squamous-cell tumors, both benign and malignant, were observed in the ear canals (Zymbal's Gland). These tumors were similar in morphology to the squamous-cell tumors observed in the skin.

Ten tumors involved the brain and nasal cavity. Seven of the tumors were types which occur very rarely in untreated Sprague-Dawley rats. These tumors were neuroepitheliomas (neuroblastomas) and nasal carcinomas, NOS.

Lymphocytic leukemias that involved multiple organs including various lymph nodes, spleen, liver, lungs, ovaries, uterus, kidneys, and bone marrow were the primary hematopoietic tumors seen. A smaller number of animals had granulocytic leukemias.

Uterine adenocarcinomas were characterized by neoplastic epithelial cells having large vesicular nuclei, prominent eosinophilic nucleoli, and ample cytoplasm that were arranged into glands. The glands were usually separated by fibrovascular stroma. The neoplastic glandular tissue arose in the endometrium and both projected into the uterine lumen and infiltrated the underlying muscle layers. Frequently the neoplastic cells penetrated the serosal surface of the uterus with transplantation on the mesentery throughout the abdominal cavity. Pulmonary metastases were also frequent. One low-dose female had a cervical uterine squamous-cell carcinoma. A few animals had poorly differentiated spindle-cell uterine sarcomas (sarcomas, NOS) and endometrial stromal polyps.

In addition to the neoplastic lesions, a number of degenerative,

proliferative, and inflammatory changes were encountered also in animals of the dosed and control groups (Appendix C). Most of these nonneoplastic lesions are commonly seen in aged Sprague-Dawley rats.

None of the high-dose male or female rats were processed for microscopic evaluation because of the short life spans, and thereby, the lack of time for tumor development. These deaths which occurred at less than 150 days were assumed to be the result of chemical toxicity. Many mid-dose males had reduced life spans that were associated with moderate to severe suppurative bronchopneumonia and bone-marrow atrophy. The lower incidence of tumors in this group may have resulted in part from the reduced life span. The female groups had fewer early deaths associated with bone-marrow atrophy and bronchopneumonias. The effects in the females of reduced life spans from toxicity were not considered important. Chronic and interstitial nephritis occurred frequently, but did not appear to be chemical related.

Administration of thio-TEPA resulted in an increase of epithelial tumors in the skin, ear canal, nasal cavity, and uterus; of hematopoietic tumors, primarily lymphocytic leukemias; of spindle-cell sarcomas in the subcutis and various other locations; and of neuroepithelial brain tumors.

Based on the histopathologic examination, thio-TEPA given intraperitoneally to Sprague-Dawley rats at doses of 0.7 and 1.4 mg/kg appeared to be carcinogenic under the conditions of this bioassay. The dose of 2.8 mg/kg proved to be too toxic for an evaluation of carcinogenic activity.

D. Statistical Analyses of Results (Rats)

Tables E1-E8 in Appendix E contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals in one group and with an incidence of at least 5% in one or more than one group. Analyses of the incidences in the high-dose groups are not included in the tables, because, as a result of the low survival, animals in these groups were not examined histopathologically. The untreated controls also are not included in the tables and in the analyses because the test conditions of the vehicle controls more closely resemble those of the dosed rats. Since the low-dose and mid-dose vehicle controls were started at different dates, the low-dose and mid-dose groups are analyzed separately with their respective controls. Due to the low survivals of animals in the dosed groups, time-adjusted analyses were performed, eliminating animals that died before week 52 on study, unless a tumor was found at the anatomic site of interest before week 52. When such an early tumor was found, comparisons were based exclusively on animals that survived at

least as long as the animal in which the first tumor was found. The statistical narrative in the following paragraphs is based on time-adjusted data only.

In rats of each sex, when the vehicle controls are used, there is no significant positive result in either the low-dose or mid-dose groups when the Bonferroni probability value of 0.025 is used as a significance level.

In male rats, when the pooled controls are used, a significant incidence of squamous-cell carcinoma in either the skin or ear canal is observed in both the low-dose and mid-dose groups ($P = 0.009$, low-dose; $P = 0.023$, mid-dose). In female rats, a significant incidence of squamous-cell carcinoma in either the skin or ear canal is observed in the mid-dose group ($P < 0.001$); however, the incidence was not significant in the low-dose group. While the vehicle-control groups were too small for effective analyses, the significant incidence of this tumor observed in both male and female rats, using pooled controls, suggests a dose association with the administration of thio-TEPA.

Leukemias were reported at a significant incidence ($P = 0.001$) in mid-dose male rats when compared with the pooled controls. When the incidences of leukemia or lymphoma are combined for analyses, a significant incidence ($P = 0.020$) is present in the low-dose

male group when compared with that in the pooled controls. The combination of leukemia or lymphoma did not appear at a significant incidence in the female groups. These results are based on the use of the pooled controls and appeared in significant incidences in males only; therefore, while suggestive of a dose relationship, the statistical evidence is not as clear as for squamous-cell carcinoma.

In female rats, adenocarcinoma was seen in a significant incidence in the mammary gland ($P = 0.006$) and in the uterus ($P = 0.001$) when the incidences in the mid-dose group were compared with those in the pooled controls. The incidence of mammary adenocarcinoma in the low-dose group also was higher than that in the pooled controls ($P = 0.033$), while the incidence of uterine carcinoma in the low-dose group was not significant. These data suggest a positive association between the administration of the test chemical and the formation of these tumors.

Significant results in the negative direction occurred in the incidences of pituitary tumors in the low-dose and mid-dose groups of female rats, and in the low-dose group of male rats.

In summary, the results of the statistical tests show that the incidences of squamous-cell carcinoma in the skin or ear canal in each sex, hematopoietic tumors in male rats, adenocarcinoma of

the mammary gland in female rats, and adenocarcinoma of the uterus in female rats are associated with the administration of thio-TEPA.

IV. RESULTS - MICE

A. Body Weights and Clinical Signs (Mice)

Mean body weights of low-dose male and female mice were only slightly lower than those of the controls throughout much of the study (figure 5). Those of the high-dose animals, particularly those of the females, were lower throughout their period of survival. Fluctuations in the growth curve may be due to mortality; as the size of a group diminishes, the mean body weight may be subject to wide variation. No other signs of toxicity were recorded.

B. Survival (Mice)

The Kaplan and Meier curves estimating the probabilities of survival for male and female mice administered thio-TEPA at the doses of this bioassay, together with those of the controls, are shown in figure 6.

For each sex, the Tarone test result is significant ($P < 0.001$), indicating that thio-TEPA decreased survival in a dose-related manner. At the high-dose, none of the males survived past week 56 and none of the females past week 43. At the low-dose, 15/35 (44%) of the males and 17/35 (49%) of the females survived to termination of the study. Survival was 7/15 (47%) among vehicle-





Figure 5. Growth Curves for Mice Treated with Thio-TEPA




Figure 6. Survival Curves for Mice Treated with Thio-TEPA

control males and 12/15 (80%) among vehicle-control females in the bioassay. Statistical analyses were time-adjusted, eliminating those mice dying before week 52; however, in some groups, insufficient numbers of mice were at risk for the development of late-appearing tumors.

C. Pathology (Mice)

Histopathologic findings on neoplasms in mice are summarized in Appendix B, tables B1 and B2; findings on nonneoplastic lesions are summarized in Appendix D, tables D1 and D2.

Excluding the lymphoreticular and epithelial tumors, the neoplasms listed in Appendix B occurred with approximately equal, or greater, frequency in the control mice than in the dosed mice, or occurred in insufficient numbers for accurate evaluation. These lesions are not uncommon in the B6C3F1 strain of mouse independent of the administration of any chemical.

Administration of thio-TEPA increased the frequency of skin tumors and lymphoreticular tumors. The distribution of these tumors was as follows:

<u>MALE</u>	<u>Vehicle Control</u>	<u>Untreated Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Number of Mice Necropsied	(14)	(15)	(30)	(34)
<u>Skin</u> (including preputial glands and subcutaneous tissues)				
Squamous-cell or adnexal carcinoma	0	0	14	1
Basal-cell carcinoma	1	0	0	0
Carcinoma, NOS	0	0	1	0
<u>Multiple Organs, Lymphoreticular</u>				
Malignant lymphoma	0	1	2	16
Leukemia, lymphocytic	1	0	0	10
Leukemia, granulocytic	0	0	1	0
<u>FEMALE</u>				
Number of Mice Necropsied	(15)	(15)	(30)	(32)
<u>Multiple Organs, Lymphoreticular</u>				
Malignant lymphoma	0	1	2	20
Leukemia, lymphocytic	0	0	3	12

The morphology of the skin tumors was similar to that described previously for the rats. The majority of the squamous-cell tumors were malignant, occurred in the groin or pelvic area, and were associated with preputial glands. Two squamous-cell carcinomas metastasized to the lungs. One mouse had a poorly differentiated spindle-cell carcinoma.

The malignant lymphomas were classified as lymphocytic, histiocytic, or mixed types and were usually solid tumors

involving multiple organs including the spleen, lymph nodes, and liver.

In addition to the malignant lymphomas, many female mice had lymphocytic leukemias. These leukemias were composed of neoplastic lymphocytes similar in appearance to those seen in the malignant lymphomas, but they had a more generalized distribution with sinusoidal involvement of the liver. One mouse had a granulocytic leukemia that was composed of well-differentiated neutrophils.

In addition to the neoplastic lesions, a number of degenerative, proliferative, and inflammatory changes were encountered also in animals of the dosed and control groups (Appendix D). These nonneoplastic lesions are commonly seen in aged B6C3F1 mice. The high-dose male and female mice had reduced life spans which were associated with the high incidence of lymphoreticular neoplasms.

Based on the histopathologic examination, thio-TEPA given intraperitoneally to B6C3F1 mice was carcinogenic under conditions of this bioassay. The tumors induced were malignant lymphomas and lymphocytic leukemias in males and females and squamous-cell carcinomas of the skin in males.

D. Statistical Analyses of Results (Mice)

Tables F1-F4 in Appendix F contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals in one group and with an incidence of at least 5% in one or more than one group. The untreated controls are not included in the tables and analyses, because the test conditions of the vehicle controls more closely resemble those of the dosed mice. Due to the low survival of the high-dose mice, time-adjusted analyses were performed, eliminating animals that died before week 52 on study, unless a tumor was found at the anatomic site of interest before week 52. When such an early tumor was found, comparisons were based exclusively on animals that survived at least as long as the animal in which the first tumor was found. The statistical narrative in the following paragraphs is based on time-adjusted data only.

In male mice, the results of the Cochran-Armitage test for dose-related trend in the incidence of squamous-cell carcinoma at all sites are significant ($P = 0.018$), and the Fisher exact test shows that the incidence in the low-dose group is significantly higher ($P = 0.004$) than that in the matched controls. There were only two animals at risk in the high-dose group, and the squamous-cell carcinoma occurred in one of these two animals. This tumor was not present in significant incidences in female

mice; however, the statistical analysis suggests a positive dose association in male mice.

The results of the Cochran-Armitage test for positive dose-related trend in the combined incidence of lymphoma or leukemia in both male and female mice are significant ($P < 0.001$). In the female high-dose group, the incidence is 32/32 (100%) and in the male high-dose group it is 26/28 (93%). The results of the Fisher exact show that the incidences in the high-dose groups are significantly higher ($P < 0.001$) than those in either type of control groups. The statistical conclusion is that the incidences of these hematopoietic tumors in mice are dose associated.

In summary, the results of the statistical tests show that the incidences of squamous-cell carcinoma at all sites in male mice and of the hematopoietic tumors in each sex are associated with the administration of thio-TEPA.

V. DISCUSSION

Under the conditions of this bioassay, thio-TEPA was toxic to both rats and mice, as evidenced by reduced mean body weights and by high mortality. Mean body weights of high-, mid-, and low-dose male rats, high- and mid-dose female rats, and high-dose male and female mice were all markedly lower than those of corresponding controls. All high-dose male rats died by week 19, all high-dose females by week 21, all mid-dose males by week 78, 91% of the mid-dose females by termination week 81, and 85% of the low-dose males and 51% of the low-dose females by termination week 82. Histopathologic evaluation of tissues was performed only on the low- and mid-dose groups of rats, since all high-dose males and females died before the occurrence of tumors could be expected. All of the high-dose male mice died by week 56, all of the high-dose females by week 43, and 77% of the low-dose males and 51% of the low-dose females by termination week 86. Because of this high mortality, time-adjusted analyses were performed on incidences of tumors in each species and each sex.

In rats, the incidence of combined neoplasms of the hematopoietic system (lymphoma, lymphocytic leukemia, or granulocytic leukemia) was significant in the males in both the low-dose ($P = 0.020$) and mid-dose ($P = 0.001$) groups, using pooled controls (pooled

controls 0/29, low-dose 6/34; pooled controls 0/30, mid-dose 6/16).

Squamous-cell carcinoma of the skin or ear canal occurred at a significant incidence in the male rats in both the low-dose ($P = 0.009$) and mid-dose ($P = 0.023$) groups, using pooled controls (pooled controls 0/29, low-dose 7/33; pooled controls 0/30, mid-dose 3/13) and in the mid-dose females ($P < 0.001$), using pooled controls (pooled controls 0/28, mid-dose 8/21); in addition, two low-dose females had such tumors, with none occurring in the corresponding low-dose controls.

The incidence of adenocarcinoma of the uterus was significant in the mid-dose female rats ($P = 0.001$), using pooled controls (pooled controls 0/28, mid-dose 7/21); in addition, two low-dose females had adenocarcinoma of the uterus, with no such tumor occurring in the corresponding low-dose controls. Adenocarcinoma of the mammary gland occurred at a significant incidence in the mid-dose female rats ($P = 0.006$), using pooled controls (pooled controls 1/28, mid-dose 8/24). In addition, seven low-dose females had these tumors, but they were not significant, since one of the low-dose pooled-control females had the tumor; and 3/10 low-dose untreated-control females had the tumor. Thus, the occurrence of mammary adenocarcinoma in the female rats cannot be clearly related to administration of the test chemical.

Also in rats, neuroepitheliomas or nasal carcinomas occurred in three low-dose males, two low-dose females, and two mid-dose females. Although these are not statistically significant incidences, these tumors did not occur among control animals and no such tumors have occurred in 380 or more Sprague-Dawley control rats of each sex in other bioassays at the same laboratory. Thus, they may be associated with administration of the chemical.

In the high-dose groups of both male and female mice, but not in the low-dose groups, the incidences of lymphoma or lymphocytic leukemia were significantly higher ($P < 0.001$) for each sex than those of either the vehicle or pooled controls (males: vehicle controls 1/8, pooled controls 1/18, low-dose 2/24, high-dose 26/28; females: vehicle controls 0/14, pooled controls 0/29, low-dose 5/26, high-dose 32/32). These tumors were observed even though all of the high-dose males died as early as week 56 and all of the high-dose females as early as week 43. In addition, granulocytic leukemia occurred in one low-dose male mouse.

In the low-dose male mice, squamous-cell carcinoma was found in the skin of seven animals, in the preputial glands of six animals, and in the ear canal of two animals. A carcinoma of the preputial gland was also found in a high-dose male. When the incidences of the tumors at the different sites were combined,

the incidence in the low-dose group, but not in the high-dose group was statistically significant using either the vehicle ($P = 0.004$) or the pooled ($P < 0.001$) controls (vehicle controls 0/8, pooled controls 0/18, low-dose 14/24, high-dose 1/2).

In previous tests of thio-TEPA for carcinogenicity, Schmähl and Osswald (1970) observed a variety of tumors in male rats of the BR 46 strain that had been injected intravenously with 1 mg/kg body weight weekly for 52 weeks. Shimkin et al. (1966), using a pulmonary tumor test system, reported the induction of lung tumors in A/J strain mice given intraperitoneal injections of thio-TEPA three times per week for 4 weeks and observed for 39 weeks after the first injection. Stoner et al. (1973), using the same test system, reported the induction of lung tumors in mice of the A/He strain given total dosages of the chemical of 4.7 or 9.4 mg/kg by intraperitoneal injection three times per week for 4 weeks and observed 24 weeks after the first injection.

It is concluded that under the conditions of this bioassay, thio-TEPA was carcinogenic in both Sprague-Dawley rats and B6C3F1 mice. In the rats, the chemical induced squamous-cell carcinoma of the skin or ear canal in both males and females, and hematopoietic neoplasms in the males; in the mice, it induced lymphoma or lymphocytic leukemia in both sexes and squamous-cell carcinoma in the skin and associated glands of males.

VI. BIBLIOGRAPHY

- Armitage, P., Statistical Methods in Medical Research, John Wiley & Sons, Inc., New York, 1971, pp. 362-365.
- Berenblum, I., ed., Carcinogenicity Testing, A Report on the Panel on Carcinogenicity of the Cancer Research Commission of the UICC, Vol. 2, International Union Against Cancer, Geneva, 1969.
- Calabresi, P. and Parks, R. E., Jr., Alkylating agents, antimetabolites, hormones, and other antiproliferative agents. In: The Pharmacological Basis of Therapeutics, Goodman, L. S., and Gilman, A., eds., Macmillan Publishing Co., Inc., New York, 1975, pp. 1254-1259 and 1265-1266.
- Calabresi, P. and Welch, A. D., Chemotherapy in neoplasms. Ann. Rev. Med. 13:147-202, 1962.
- Carter, S. K. and Slavik, M., Chemotherapy of cancer. Ann Rev. Pharmacol. 14:157, 1974.
- Cox, D. R., Regression models and life tables. J. R. Statist. Soc. B 34 (2):187-220, 1972.
- Cox, D. R., Analysis of Binary Data, Methuen & Co., Ltd., London, 1970, pp. 48-52.
- Gart, J. J., The comparison of proportions: a review of significance tests, confidence limits and adjustments for stratification. Rev. Int. Statist. Inst. 39:148-169, 1971.
- Kaplan, E. L. and Meier, P., Nonparametric estimation from incomplete observations. J. Amer. Statist. Assoc. 53:457-481, 1958.
- Lederle Laboratories. Physicians' Desk Reference, Baker, C. E., Jr., Medical Economics Co., Oradell, N. J., 1977, pp. 903 and 906.
- Linhart, M. S., Cooper, J., Martin, R. L., Page, N., and Peters, J., Carcinogenesis bioassay data system. Comp. and Biomed. Res. 7:230-248, 1974.

- Miller, R. G., Jr., Simultaneous Statistical Inference, McGraw-Hill Book Co., New York, 1966, pp. 6-10.
- Montgomery, J. A. and Struck, R. J., The relationship of the metabolism of anticancer agents to their activity. Progress in Drug Research 17:363-364, 1973.
- Saffiotti, U., Montesano, R., Sellakumar, A. R., Cefis, F., and Kaufman, D. G., Respiratory tract carcinogenesis in hamsters induced by different numbers of administrations of benzo(a) pyrene and ferric oxide. Cancer Res. 32:1073-1081, 1972.
- Schmähl, V. D. and Osswald, H., Experimentelle Untersuchungen über carcinogene Wirkungen von Krebs-Chemotherapeutica und Immunosuppressiva. Arzneimittel-Forschung 20 (10):1461-1470, 1970.
- Shimkin, M. B., Weisburger, J. H., Weisburger, E. K., Guboreff, N., and Suntzeff, V., Bioassay of 29 alkylating chemicals by the pulmonary-tumor response in strain A mice. J. Natl. Cancer Inst. 36:915-935, 1966.
- Stoner, G. D., Shimkin, M. B., Kniazeff, A. J., Weisburger, J. H., Weisburger, E. K., and Gori, G. B., Test for carcinogenicity of food additives and chemotherapeutic agents by the pulmonary tumors response in strain A mice. Cancer Res. 33:3069-3085, 1973.
- Tarone, R. E., Tests for trend in life table analysis. Biometrika 62(3):679-682, 1975.
- U.S. Patent 2,670,347 (February 23, 1954).
- Wheeler, G. P., Alkylating agents. In: Cancer Medicine, Holland, J. F., and Frei, E., III, eds., Lea & Febiger, Philadelphia, 1973, pp. 791-805.

APPENDIX A

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA

TABLE A1.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (CONTROL GROUPS)**

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
ANIMALS INITIALLY IN STUDY	10	10	10	10
ANIMALS NECROPSIED	10	9	10	10
ANIMALS EXAMINED HISTOPATHOLOGICALLY	10	9	10	10
INTEGUMENTARY SYSTEM				
*SKIN	(10)	(9)	(10)	(10)
SQUAMOUS CELL PAPILLOMA				1 (10%)
FIBROMA			1 (10%)	
*SUBCUT TISSUE	(10)	(9)	(10)	(10)
FIBROMA			1 (10%)	
RESPIRATORY SYSTEM				
NONE				
HEMATOPOIETIC SYSTEM				
NONE				
CIRCULATORY SYSTEM				
NONE				
DIGESTIVE SYSTEM				
NONE				
URINARY SYSTEM				
NONE				
ENDOCRINE SYSTEM				
*PITUITARY	(10)	(7)	(10)	(9)
CHROMOPHOBE ADENOMA		1 (14%)		3 (33%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE A1. MALE RATS (CONTROL GROUPS): NEOPLASMS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
CHROMOPHOBE CARCINOMA				1 (11%)
REPRODUCTIVE SYSTEM				
*MAMMARY GLAND FIBROADENOMA	(10) 2 (20%)	(9)	(10) 1 (10%)	(10)
NERVOUS SYSTEM				
NONE				
SPECIAL SENSE ORGANS				
NONE				
MUSCULOSKELETAL SYSTEM				
NONE				
BODY CAVITIES				
NONE				
ALL OTHER SYSTEMS				
NONE				
ANIMAL DISPOSITION SUMMARY				
ANIMALS INITIALLY IN STUDY	10	10	10	10
NATURAL DEATH@	2	2	1	
MORIBUND SACRIFICE			1	
SCHEDULED SACRIFICE				
ACCIDENTALLY KILLED				
TERMINAL SACRIFICE	8	8	8	10
ANIMAL MISSING				
@ INCLUDES AUTOLYZED ANIMALS				
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE A1. MALE RATS (CONTROL GROUPS): NEOPLASMS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
TUMOR SUMMARY				
TOTAL ANIMALS WITH PRIMARY TUMORS*	2	1	3	5
TOTAL PRIMARY TUMORS	2	1	3	5
TOTAL ANIMALS WITH BENIGN TUMORS	2	1	3	4
TOTAL BENIGN TUMORS	2	1	3	4
TOTAL ANIMALS WITH MALIGNANT TUMORS				1
TOTAL MALIGNANT TUMORS				1
TOTAL ANIMALS WITH SECONDARY TUMORS#				
TOTAL SECONDARY TUMORS				
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT				
TOTAL UNCERTAIN TUMORS				
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC				
TOTAL UNCERTAIN TUMORS				
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS				
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN				

TABLE A2.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (TREATED GROUPS)**

	LOW DOSE	MID DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	39	35	35
ANIMALS NECROPSIED	37	30	32
ANIMALS EXAMINED HISTOPATHOLOGICALLY	37	29	0
INTEGUMENTARY SYSTEM			
*SKIN	(37)	(30)	(32)
SQUAMOUS CELL PAPILLOMA	1 (3%)		
SQUAMOUS CELL CARCINOMA	5 (14%)	3 (10%)	
BASAL-CELL CARCINOMA		1 (3%)	
TRICHOEPITHELIOMA	1 (3%)		
*SUBCUT TISSUE	(37)	(30)	(32)
SARCOMA, NOS	5 (14%)		
FIBROMA	2 (5%)		
FIBROSARCOMA	1 (3%)		
RESPIRATORY SYSTEM			
*NASAL CAVITY	(37)	(30)	(32)
CARCINOMA, NOS	1 (3%)		
#LUNG	(36)	(29)	
SQUAMOUS CELL CARCINOMA, METASTA		1 (3%)	
ALVEOLAR/BRONCHIOLAR ADENOMA		1 (3%)	
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(37)	(30)	(32)
MALIG. LYMPHOMA, HISTIOCYTIC TYPE	1 (3%)		
LYMPHOCYTIC LEUKEMIA	3 (8%)	5 (17%)	
GRANULOCYTIC LEUKEMIA	2 (5%)	1 (3%)	
CIRCULATORY SYSTEM			
<u>NONE</u>			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2. MALE RATS (TREATED GROUPS): NEOPLASMS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
DIGESTIVE SYSTEM			
*LIVER CARCINOMA, NOS	(37) 1 (3%)	(29) 1 (3%)	
URINARY SYSTEM			
*KIDNEY HAMARTOMA	(37) 1 (3%)	(29)	
ENDOCRINE SYSTEM			
*PITUITARY CHROMOPHOBE ADENOMA	(28) 1 (4%)	(24) 1 (4%)	
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND ADENOCARCINOMA, NOS	(37) 1 (3%)	(30)	(32)
NERVOUS SYSTEM			
*EPRAIN SARCOMA, NOS	(36)	(28) 1 (4%)	
ASTROCYTOMA	1 (3%)		
NEUROBLASTOMA	1 (3%)		
OLFACTORY NEUROELASTOMA	1 (3%)		
SPECIAL SENSE ORGANS			
*EAR CANAL SQUAMOUS CELL PAPILLOMA	(37) 1 (3%)	(30)	(32)
SQUAMOUS CELL CARCINOMA	3 (8%)	1 (3%)	
MUSCULOSKELETAL SYSTEM			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2. MALE RATS (TREATED GROUPS): NEOPLASMS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
BODY CAVITIES			
*MEDIASTINUM SARCOMA, NOS	(37) 1 (3%)	(30)	(32)
*ABDOMINAL CAVITY SARCOMA, NOS	(37)	(30) 1 (3%)	(32)
*MESENTERY SARCOMA, NOS	(37) 2 (5%)	(30)	(32)
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS SQUAMOUS CELL CARCINOMA, METASTA	(37)	(30) 1 (3%)	(32)
CRANIAL CAVITY SARCOMA, NOS	1		
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	39	35	35
NATURAL DEATH@	7	13	26
MORIBUND SACRIFICE	26	22	9
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	6		
ANIMAL MISSING			
@ INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2. MALE RATS (TREATED GROUPS): NEOPLASMS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	27	14	
TOTAL PRIMARY TUMORS	36	16	
TOTAL ANIMALS WITH BENIGN TUMORS	6	2	
TOTAL BENIGN TUMORS	7	2	
TOTAL ANIMALS WITH MALIGNANT TUMORS	26	12	
TOTAL MALIGNANT TUMORS	29	14	
TOTAL ANIMALS WITH SECONDARY TUMORS#		2	
TOTAL SECONDARY TUMORS		2	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE A3.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (CONTROL GROUPS)**

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
ANIMALS INITIALLY IN STUDY	10	10	10	10
ANIMALS NECROPSIED	10	10	10	10
ANIMALS EXAMINED HISTOPATHOLOGICALLY	10	10	10	10
INTEGUMENTARY SYSTEM				
*SUBCUT TISSUE FIBROMA	(10) 1 (10%)	(10)	(10)	(10)
RESPIRATORY SYSTEM				
NONE				
HEMATOPOIETIC SYSTEM				
NONE				
CIRCULATORY SYSTEM				
NONE				
DIGESTIVE SYSTEM				
NONE				
URINARY SYSTEM				
NONE				
ENDOCRINE SYSTEM				
#PITUITARY CHROMOPHOBE ADENOMA	(10) 5 (50%)	(10) 6 (60%)	(10) 2 (20%)	(8) 1 (13%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE A3. FEMALE RATS (CONTROL GROUPS): NEOPLASMS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
CHROMOPHOBE CARCINOMA		1 (10%)		
REPRODUCTIVE SYSTEM				
*MAMMARY GLAND	(10)	(10)	(10)	(10)
ADENOCARCINOMA, NOS		3 (30%)	1 (10%)	
FIBROADENOMA	2 (20%)	6 (60%)	5 (50%)	2 (20%)
NERVOUS SYSTEM				
NONE				
SPECIAL SENSE ORGANS				
NONE				
MUSCULOSKELETAL SYSTEM				
NONE				
BODY CAVITIES				
NONE				
ALL OTHER SYSTEMS				
NONE				
ANIMAL DISPOSITION SUMMARY				
ANIMALS INITIALLY IN STUDY	10	10	10	10
NATURAL DEATH@	2			1
MORIBUND SACRIFICE		2		
SCHEDULED SACRIFICE				
ACCIDENTALLY KILLED		1		
TERMINAL SACRIFICE	8	7	10	9
ANIMAL MISSING				
@ INCLUDES AUTOLYZED ANIMALS				
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE A3. FEMALE RATS (CONTROL GROUPS): NEOPLASMS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
TUMOR SUMMARY				
TOTAL ANIMALS WITH PRIMARY TUMORS*	7	9	5	3
TOTAL PRIMARY TUMORS	8	16	8	3
TOTAL ANIMALS WITH BENIGN TUMORS	7	9	5	3
TOTAL BENIGN TUMORS	8	12	7	3
TOTAL ANIMALS WITH MALIGNANT TUMORS		4	1	
TOTAL MALIGNANT TUMORS		4	1	
TOTAL ANIMALS WITH SECONDARY TUMORS#				
TOTAL SECONDARY TUMORS				
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT				
TOTAL UNCERTAIN TUMORS				
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC				
TOTAL UNCERTAIN TUMORS				
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS				
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN				

TABLE A4.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (TREATED GROUPS)**

	LOW DOSE	MID DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	31	35	35
ANIMALS NECROPSIED	30	33	34
ANIMALS EXAMINED HISTOPATHOLOGICALLY	30	33	0
INTEGUMENTARY SYSTEM			
*SKIN	(30)	(33)	(34)
SQUAMOUS CELL PAPILLOMA		1 (3%)	
SQUAMOUS CELL CARCINOMA		3 (9%)	
*SUBCUT TISSUE	(30)	(33)	(34)
SARCOMA, NOS	1 (3%)		
FIBROMA	1 (3%)		
FIBROSARCOMA	1 (3%)		
RESPIRATORY SYSTEM			
*NASAL CAVITY	(30)	(33)	(34)
CARCINOMA, NOS		1 (3%)	
*LUNG	(30)	(33)	
ADENOCARCINOMA, NOS, METASTATIC	2 (7%)		
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(30)	(33)	(34)
MALIG. LYMPHOMA, HISTIOCYTIC TYPE		1 (3%)	
LYMPHOCYTIC LEUKEMIA		1 (3%)	
GRANULOCYTIC LEUKEMIA	1 (3%)		
*MANDIBULAR L. NODE	(28)	(1)	
MALIG. LYMPHOMA, UNDIFFER-TYPE	1 (4%)		
*THYMUS	(30)	(33)	
MALIG. LYMPHOMA, LYMPHOCYTIC TYPE		1 (3%)	
CIRCULATORY SYSTEM			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A4. FEMALE RATS (TREATED GROUPS): NEOPLASMS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
DIGESTIVE SYSTEM			
#SALIVARY GLAND SARCOMA, NOS	(24) 1 (4%)		
URINARY SYSTEM			
#KIDNEY HAMARTOMA	(30) 1 (3%)	(33)	
ENDOCRINE SYSTEM			
#PITUITARY CHROMOPHOBE ADENOMA	(29) 4 (14%)	(32)	
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(30)	(33)	(34)
ADENOMA, NOS		3 (9%)	
ADENOCARCINOMA, NOS	7 (23%)	8 (24%)	
FIBROADENOMA	8 (27%)	4 (12%)	
#UTERUS	(30)	(32)	
ADENOCARCINOMA, NOS	2 (7%)	7 (22%)	
SARCOMA, NOS	2 (7%)	1 (3%)	
ENDOMETRIAL STROMAL POLYP	4 (13%)	1 (3%)	
#CERVIX UTERI	(30)	(32)	
SQUAMOUS CELL CARCINOMA	1 (3%)		
#OVARY	(30)	(32)	
LUTEOMA		1 (3%)	
SARCOMA, NOS		1 (3%)	
NERVOUS SYSTEM			
#BRAIN	(30)	(32)	
SARCOMA, NOS		1 (3%)	
OLFACTORY NEUROBLASTOMA	2 (7%)	1 (3%)	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A4. FEMALE RATS (TREATED GROUPS): NEOPLASMS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
SPECIAL SENSE ORGANS			
*EAR CANAL	(30)	(33)	(34)
SQUAMOUS CELL PAPILLOMA		1 (3%)	
SQUAMOUS CELL CARCINOMA	2 (7%)	5 (15%)	
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY	(30)	(33)	(34)
ADENOCARCINOMA, NOS, METASTATIC	1 (3%)	1 (3%)	
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(30)	(33)	(34)
ADENOCARCINOMA, NOS		1 (3%)	
ADENOCARCINOMA, NOS, METASTATIC		4 (12%)	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	31	35	35
NATURAL DEATH ^a	9	10	15
MORIBUND SACRIFICE	9	22	20
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	13	3	
ANIMAL MISSING			
^a INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A4. FEMALE RATS (TREATED GROUPS): NEOPLASMS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	26	26	
TOTAL PRIMARY TUMORS	39	43	
TOTAL ANIMALS WITH BENIGN TUMORS	14	9	
TOTAL BENIGN TUMORS	18	11	
TOTAL ANIMALS WITH MALIGNANT TUMORS	17	22	
TOTAL MALIGNANT TUMORS	21	32	
TOTAL ANIMALS WITH SECONDARY TUMORS*	3	5	
TOTAL SECONDARY TUMORS	3	5	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX B

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MICE
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA

TABLE B1.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MICE
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA**

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	15	15	@34	35
ANIMALS NECROPSIED	14	15	30	34
ANIMALS EXAMINED HISTOPATHOLOGICALLY	14	15	30	33
INTEGUMENTARY SYSTEM				
*SKIN	(14)	(15)	(30)	(34)
CARCINOMA, NOS			1 (3%)	
SQUAMOUS CELL CARCINOMA			7 (23%)	
BASAL-CELL CARCINOMA	1 (7%)			
*SUBCUT TISSUE	(14)	(15)	(30)	(34)
SQUAMOUS CELL CARCINOMA			1 (3%)	
SARCOMA, NOS			1 (3%)	
RESPIRATORY SYSTEM				
*LUNG	(14)	(15)	(30)	(33)
SQUAMOUS CELL CARCINOMA, METASTA			1 (3%)	1 (3%)
ALVEOLAR/BRONCHIOLAR ADENOMA	1 (7%)	4 (27%)	5 (17%)	
HEMATOPOIETIC SYSTEM				
*MULTIPLE ORGANS	(14)	(15)	(30)	(34)
MALIGNANT LYMPHOMA, NOS				2 (6%)
MALIG. LYMPHOMA, LYMPHOCYTIC TYPE				14 (41%)
MALIG. LYMPHOMA, HISTIOCYTIC TYPE			1 (3%)	
MALIGNANT LYMPHOMA, MIXED TYPE			1 (3%)	
LYMPHOCYTIC LEUKEMIA	1 (7%)			10 (29%)
GRANULOCYTIC LEUKEMIA			1 (3%)	
*MESENTERIC L. NODE	(2)	(5)	(1)	(15)
MALIGNANT LYMPHOMA, NOS		1 (20%)		
CIRCULATORY SYSTEM				
*MYOCARDIUM	(14)	(15)	(30)	(32)
HEMANGIOMA			1 (3%)	

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

@NOTE: 35 ANIMALS WERE INITIALLY IN STUDY BUT ONE WAS FOUND TO BE A FEMALE IN A MALE GROUP AND WAS DELETED.

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
DIGESTIVE SYSTEM				
#LIVER	(14)	(15)	(30)	(33)
HEPATOCELLULAR ADENOMA	1 (7%)	2 (13%)	3 (10%)	
HEPATOCELLULAR CARCINOMA	1 (7%)		1 (3%)	
URINARY SYSTEM				
NONE				
ENDOCRINE SYSTEM				
NONE				
REPRODUCTIVE SYSTEM				
*PREPUTIAL GLAND	(14)	(15)	(30)	(34)
SQUAMOUS CELL CARCINOMA			6 (20%)	1 (3%)
NERVOUS SYSTEM				
NONE				
SPECIAL SENSE ORGANS				
*EAR CANAL	(14)	(15)	(30)	(34)
SQUAMOUS CELL PAPILLOMA			1 (3%)	
SQUAMOUS CELL CARCINOMA			2 (7%)	
MUSCULOSKELETAL SYSTEM				
NONE				
BODY CAVITIES				
NONE				
ALL OTHER SYSTEMS				
NONE				
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ANIMAL DISPOSITION SUMMARY				
ANIMALS INITIALLY IN STUDY	15	15	35	35
NATURAL DEATH ^a	7	2	13	13
MORIBUND SACRIFICE			6	22
SCHEDULED SACRIFICE				
ACCIDENTALLY KILLED	1			
TERMINAL SACRIFICE	7	13	15	
ANIMAL MISSING				
ANIMAL DELETED (WRONG SEX)			1	
^a INCLUDES AUTOLYZED ANIMALS				
TUMOR SUMMARY				
TOTAL ANIMALS WITH PRIMARY TUMORS*	4	6	19	27
TOTAL PRIMARY TUMORS	5	7	32	27
TOTAL ANIMALS WITH BENIGN TUMORS	2	5	8	
TOTAL BENIGN TUMORS	2	6	10	
TOTAL ANIMALS WITH MALIGNANT TUMORS	3	1	18	27
TOTAL MALIGNANT TUMORS	3	1	22	27
TOTAL ANIMALS WITH SECONDARY TUMORS [#]			1	1
TOTAL SECONDARY TUMORS			1	1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT				
TOTAL UNCERTAIN TUMORS				
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC				
TOTAL UNCERTAIN TUMORS				
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS				
[#] SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN				

TABLE B2.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MICE
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA**

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	15	15	35	35
ANIMALS MISSING			1	2
ANIMALS NECROPSIED	15	15	30	32
ANIMALS EXAMINED HISTOPATHOLOGICALLY	15	15	30	32
INTEGUMENTARY SYSTEM				
*SKIN	(15)	(15)	(30)	(32)
SQUAMOUS CELL PAPILLOMA			1 (3%)	
RESPIRATORY SYSTEM				
#LUNG	(15)	(15)	(29)	(32)
ADENOCARCINOMA, NOS, METASTATIC			1 (3%)	
ALVEOLAR/BRONCHIOLAR ADENOMA			5 (17%)	
HEMATOPOIETIC SYSTEM				
*MULTIPLE ORGANS	(15)	(15)	(30)	(32)
MALIGNANT LYMPHOMA, NOS				1 (3%)
MALIG. LYMPHOMA, LYMPHOCYTIC TYPE			2 (7%)	19 (59%)
MALIGNANT LYMPHOMA, MIXED TYPE		1 (7%)		
LYMPHOCYTIC LEUKEMIA			3 (10%)	12 (38%)
CIRCULATORY SYSTEM				
NONE				
DIGESTIVE SYSTEM				
#LIVER	(15)	(15)	(30)	(32)
HEPATOCELLULAR ADENOMA			2 (7%)	
URINARY SYSTEM				
NONE				
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ENDOCRINE SYSTEM				
*ADRENAL	(15)	(15)	(28)	(32)
CORTICAL CARCINOMA			1 (4%)	
PHEOCHROMOCYTOMA			1 (4%)	
REPRODUCTIVE SYSTEM				
*MAMMARY GLAND	(15)	(15)	(30)	(32)
ADENOCARCINOMA, NOS			1 (3%)	
*OVARY	(15)	(15)	(29)	(31)
CARCINOMA, NOS			3 (10%)	
NERVOUS SYSTEM				
NONE				
SPECIAL SENSE ORGANS				
*EAR CANAL	(15)	(15)	(30)	(32)
SQUAMOUS CELL PAPILLOMA			1 (3%)	
MUSCULOSKELETAL SYSTEM				
NONE				
BODY CAVITIES				
NONE				
ALL OTHER SYSTEMS				
NONE				
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ANIMAL DISPOSITION SUMMARY				
ANIMALS INITIALLY IN STUDY	15	15	35	35
NATURAL DEATH ^a	2	1	11	16
MORIBUND SACRIFICE	1		4	17
SCHEDULED SACRIFICE				
ACCIDENTALLY KILLED			2	
TERMINAL SACRIFICE	12	14	17	
ANIMAL MISSING			1	2
^a INCLUDES AUTOLYZED ANIMALS				
TUMOR SUMMARY				
TOTAL ANIMALS WITH PRIMARY TUMORS*		1	17	32
TOTAL PRIMARY TUMORS		1	20	32
TOTAL ANIMALS WITH BENIGN TUMORS			10	
TOTAL BENIGN TUMORS			10	
TOTAL ANIMALS WITH MALIGNANT TUMORS		1	9	32
TOTAL MALIGNANT TUMORS		1	10	32
TOTAL ANIMALS WITH SECONDARY TUMORS [#]			1	
TOTAL SECONDARY TUMORS			1	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT				
TOTAL UNCERTAIN TUMORS				
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC				
TOTAL UNCERTAIN TUMORS				
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS				
[#] SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN				

APPENDIX C

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA

11/11/11

TABLE C1.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (CONTROL GROUPS)**

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
ANIMALS INITIALLY IN STUDY	10	10	10	10
ANIMALS NECROPSIED	10	9	10	10
ANIMALS EXAMINED HISTOPATHOLOGICALLY	10	9	10	10
INTEGUMENTARY SYSTEM				
*SKIN	(10)	(9)	(10)	(10)
ULCER, CHRONIC				1 (10%)
RESPIRATORY SYSTEM				
#TRACHEA	(10)	(9)	(10)	(10)
INFLAMMATION, NOS	1 (10%)			
INFLAMMATION, ACUTE/CHRONIC			1 (10%)	
INFLAMMATION, CHRONIC		1 (11%)		
INFLAMMATION, CHRONIC SUPPURATIVE				1 (10%)
#LUNG/BRONCHUS	(10)	(9)	(10)	(10)
BRONCHIECTASIS		1 (11%)		
#LUNG/BRONCHIOLE	(10)	(9)	(10)	(10)
HYPERPLASIA, LYMPHOID	1 (10%)		1 (10%)	
#LUNG	(10)	(9)	(10)	(10)
BRONCHOPNEUMONIA SUPPURATIVE			1 (10%)	
HEMATOPOIETIC SYSTEM				
#BONE MARROW	(10)	(9)	(10)	(10)
ATROPHY, NOS	2 (20%)	4 (44%)	5 (50%)	7 (70%)
CIRCULATORY SYSTEM				
NONE				
DIGESTIVE SYSTEM				
#LIVER	(10)	(9)	(10)	(10)
HYPERPLASIA, NODULAR		1 (11%)		

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE C1. MALE RATS (CONTROL GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
*BILE DUCT HYPERPLASIA, NOS	(10) 1 (10%)	(9)	(10)	(10)
URINARY SYSTEM				
*KIDNEY	(10)	(9)	(10)	(10)
INFLAMMATION, INTERSTITIAL	6 (60%)		4 (40%)	
INFLAMMATION, SUPPURATIVE	1 (10%)			
INFLAMMATION, CHRONIC		6 (67%)		8 (80%)
ENDOCRINE SYSTEM				
NONE				
REPRODUCTIVE SYSTEM				
*MAMMARY GLAND CYST, NOS	(10)	(9)	(10)	(10) 1 (10%)
*PROSTATE INFLAMMATION, CHRONIC SUPPURATIV	(10)	(9) 1 (11%)	(10)	(10) 1 (10%)
*TESTIS ATROPHY, NOS	(9)	(9) 1 (11%)	(10)	(10)
NERVOUS SYSTEM				
NONE				
SPECIAL SENSE ORGANS				
NONE				
MUSCULOSKELETAL SYSTEM				
NONE				
BODY CAVITIES				
NONE				
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE C1. MALE RATS (CONTROL GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
ALL OTHER SYSTEMS				
NONE				
SPECIAL MORPHOLOGY SUMMARY				
NO LESION REPORTED		2	2	
NO NECROPSY PERFORMED		1		
• NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
• NUMBER OF ANIMALS NECROPSIED				

TABLE C2.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (TREATED GROUPS)**

	LOW DOSE	MID DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	39	35	35
ANIMALS NECROPSIED	37	30	32
ANIMALS EXAMINED HISTOPATHOLOGICALLY	37	29	0
INTEGUMENTARY SYSTEM			
*SKIN	(37)	(30)	(32)
EPIDERMAL INCLUSION CYST	2 (5%)		
DERMAL INCLUSION CYST		1 (3%)	
INFLAMMATION, SUPPURATIVE	1 (3%)		
ABSCESS, CHRONIC	1 (3%)		
HYPERKERATOSIS	1 (3%)		
KERATIN-PEARL FORMATION	1 (3%)	1 (3%)	
RESPIRATORY SYSTEM			
#TRACHEA	(37)	(28)	
INFLAMMATION, CHRONIC	2 (5%)		
#LUNG/BRONCHIOLE	(36)	(29)	
HYPERPLASIA, PLASMA CELL		2 (7%)	
#LUNG	(36)	(29)	
HEMORRHAGE	1 (3%)		
INFLAMMATION, SUPPURATIVE	1 (3%)		
BRONCHOPNEUMONIA SUPPURATIVE	2 (6%)	10 (34%)	
BRONCHOPNEUMONIA CHRONIC SUPPURA	3 (8%)	5 (17%)	
ABSCESS, CHRONIC		1 (3%)	
METAPLASIA, SQUAMOUS	1 (3%)	2 (7%)	
HYPERPLASIA, PLASMA CELL		1 (3%)	
HEMATOPOIETIC SYSTEM			
#BONE MARROW	(35)	(29)	
ATROPHY, NOS	17 (43%)	10 (34%)	
#SPLEEN	(37)	(28)	
INFLAMMATION, NECROTIZING	1 (3%)		

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE C2. MALE RATS (TREATED GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
ATROPHY, NOS	1 (3%)		
ANGIECTASIS	1 (3%)		
HEMATOPOIESIS	4 (11%)	1 (4%)	
*MANDIBULAR L. NODE	(31)	(6)	
CONGESTION, NOS	1 (3%)		
HYPERPLASIA, RETICULUM CELL	1 (3%)		
*MESENTERIC L. NODE	(31)	(6)	
CONGESTION, NOS	1 (3%)		
INFLAMMATION, GRANULOMATOUS	1 (3%)		
CYTOMEGALY	1 (3%)		
CIRCULATORY SYSTEM			
*MYOCARDIUM	(37)	(28)	
INFLAMMATION, INTERSTITIAL	1 (3%)		
CALCIFICATION, METASTATIC	1 (3%)		
DIGESTIVE SYSTEM			
*LIVER	(37)	(29)	
HEMORRHAGE	1 (3%)		
ANGIECTASIS		1 (3%)	
*PANCREAS	(37)	(29)	
NECROSIS, FAT	1 (3%)		
*ESOPHAGUS	(36)	(27)	
ULCER, NOS	1 (3%)		
INFLAMMATION, SUPPURATIVE	1 (3%)		
*GASTRIC MUCOSA	(37)	(29)	
CALCIFICATION, METASTATIC	1 (3%)		
*COLON	(36)	(29)	
HEMORRHAGE	1 (3%)		
*CECUM	(36)	(29)	
THROMBOSIS, NOS		1 (3%)	
URINARY SYSTEM			
*KIDNEY	(37)	(29)	
CALCULUS, NOS		1 (3%)	

* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE C2. MALE RATS (TREATED GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
INFLAMMATION, INTERSTITIAL		7 (24%)	
INFLAMMATION, CHRONIC	26 (70%)		
CALCIFICATION, METASTATIC	1 (3%)		
ENDOCRINE SYSTEM			
*ADRENAL	(37)	(29)	
ANGIECTASIS	1 (3%)		
REPRODUCTIVE SYSTEM			
*TESTIS	(36)	(27)	
ATROPHY, NOS	15 (42%)		
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*EYE	(37)	(30)	(32)
HEMORRHAGE	1 (3%)		
INFLAMMATION, SUPPURATIVE	1 (3%)		
*EYE/CORNEA	(37)	(30)	(32)
ULCER, CHRONIC	1 (3%)		
*EYE/CONJUNCTIVA	(37)	(30)	(32)
INFLAMMATION, CHRONIC SUPPURATIV		1 (3%)	
*MIDDLE EAR	(37)	(30)	(32)
INFLAMMATION, SUPPURATIVE	1 (3%)		
INFLAMMATION, CHRONIC SUPPURATIV	1 (3%)		
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*PERITONEUM	(37)	(30)	(32)
INFLAMMATION, CHRONIC	1 (3%)		
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2. MALE RATS (TREATED GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
*PLEURA INFLAMMATION, SUPPURATIVE	(37) 1 (3%)	(30)	(32)
*MESENTERY NECROSIS, FAT	(37) 1 (3%)	(30)	(32)
ALL OTHER SYSTEMS			
ADIPOSE TISSUE INFLAMMATION, GRANULOMATOUS	1		
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED		1	
NECROPSY PERF/NO HISTO PERFORMED		1	31
NO NECROPSY PERFORMED	1		
AUTO/NECROPSY/NO HISTO			1
AUTOLYSIS/NO NECROPSY	1	5	3
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY * NUMBER OF ANIMALS NECROPSIED			

TABLE C3.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (CONTROL GROUPS)**

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
ANIMALS INITIALLY IN STUDY	10	10	10	10
ANIMALS NECROPSIED	10	10	10	10
ANIMALS EXAMINED HISTOPATHOLOGICALLY	10	10	10	10
INTEGUMENTARY SYSTEM				
NONE				
RESPIRATORY SYSTEM				
#TRACHEA	(10)	(10)	(10)	(10)
INFLAMMATION, NOS	1 (10%)			
INFLAMMATION, CHRONIC				1 (10%)
#LUNG	(10)	(10)	(10)	(10)
PNEUMONIA, LIPID		1 (10%)		
PNEUMONIA INTERSTITIAL CHRONIC		1 (10%)		
BRONCHOPNEUMONIA CHRONIC SUPPURA		1 (10%)		
HEMATOPOIETIC SYSTEM				
#BONE MARROW	(10)	(10)	(10)	(9)
ATROPHY, NOS	4 (40%)	7 (70%)	7 (70%)	7 (78%)
CIRCULATORY SYSTEM				
NONE				
DIGESTIVE SYSTEM				
#LIVER/PERI PORTAL	(10)	(10)	(10)	(10)
FIBROSIS	1 (10%)			
#PANCREATIC ACINUS	(10)	(10)	(10)	(10)
ATROPHY, NOS	1 (10%)			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE C3. FEMALE RATS (CONTROL GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
URINARY SYSTEM				
*KIDNEY	(10)	(10)	(10)	(10)
INFLAMMATION, INTERSTITIAL			1 (10%)	
INFLAMMATION, CHRONIC		3 (30%)		1 (10%)
GLOMERULONEPHRITIS, CHRONIC	1 (10%)			
ENDOCRINE SYSTEM				
*ADRENAL	(10)	(10)	(10)	(10)
ANGIECTASIS	2 (20%)	1 (10%)	1 (10%)	3 (30%)
REPRODUCTIVE SYSTEM				
*MAMMARY GLAND	(10)	(10)	(10)	(10)
CYST, NOS	2 (20%)	1 (10%)	5 (50%)	1 (10%)
HEMORRHAGIC CYST		1 (10%)		
*UTERUS	(10)	(10)	(10)	(10)
INFLAMMATION, SUPPURATIVE			1 (10%)	
*UTERUS/ENDOMETRIUM	(10)	(10)	(10)	(10)
INFLAMMATION, SUPPURATIVE	5 (50%)	1 (10%)		
*OVARY	(10)	(10)	(9)	(10)
CYST, NOS		1 (10%)		6 (60%)
INFLAMMATION, HEMORRHAGIC		1 (10%)		
INFLAMMATION, CHRONIC SUPPURATIVE		1 (10%)		
NERVOUS SYSTEM				
*BRAIN	(10)	(10)	(10)	(9)
HYDROCEPHALUS, NCS			1 (10%)	
SPECIAL SENSE ORGANS				
NONE				
MUSCULOSKELETAL SYSTEM				
NONE				
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE C3. FEMALE RATS (CONTROL GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	MID AND HIGH DOSE UNTREATED CONTROL	LOW DOSE UNTREATED CONTROL	MID AND HIGH DOSE VEHICLE CONTROL	LOW DOSE VEHICLE CONTROL
BODY CAVITIES				
*MESENTERY STEATITIS	(10)	(10)	(10)	(10) 1 (10%)
ALL OTHER SYSTEMS				
ADIPOSE TISSUE INFLAMMATION, FOCAL GRANULOMATOU		1		
SPECIAL MORPHOLOGY SUMMARY				
NO LESION REPORTED	1		2	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE C4.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA (TREATED GROUPS)**

	LOW DOSE	MID DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	31	35	35
ANIMALS NECROPSIED	30	33	34
ANIMALS EXAMINED HISTOPATHOLOGICALLY	30	33	0
INTEGUMENTARY SYSTEM			
*SKIN	(30)	(33)	(34)
INFLAMMATION, SUPPURATIVE		1 (3%)	
HYPERKERATOSIS		1 (3%)	
PARAKERATOSIS		1 (3%)	
RESPIRATORY SYSTEM			
*TRACHEA	(30)	(33)	
INFLAMMATION, NOS		2 (6%)	
LYMPHOCYTIC INFLAMMATORY INFILTR		1 (3%)	
INFLAMMATION, CHRONIC	3 (10%)		
*LUNG/BRONCHIOLE	(30)	(33)	
METAPLASIA, SQUAMOUS		1 (3%)	
*LUNG	(30)	(33)	
EDEMA, NOS	1 (3%)		
HEMORRHAGE	1 (3%)		
BRONCHOPNEUMONIA SUPPURATIVE	1 (3%)	4 (12%)	
ABSCCESS, NOS		2 (6%)	
BRONCHOPNEUMONIA CHRONIC SUPPURA	1 (3%)	4 (12%)	
INFLAMMATION, GRANULOMATOUS	3 (10%)		
CYTOMEGALY	2 (7%)		
HEMATOPOIETIC SYSTEM			
*BONE MARROW	(29)	(33)	
ATROPHY, NOS	13 (45%)	8 (24%)	
*SPLEEN	(30)	(33)	
HEMORRHAGE	1 (3%)		
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY * NUMBER OF ANIMALS NECROPSIED			

TABLE C4. FEMALE RATS (TREATED GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
ANGIECTASIS	1 (3%)		
HEMATOPOIESIS	5 (17%)	7 (21%)	
*MESENTERIC L. NODE	(28)	(1)	
CONGESTION, NOS	2 (7%)		
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
*LIVER	(30)	(33)	
HEMORRHAGE	2 (7%)		
INFLAMMATION, CHRONIC NECROTIZING	1 (3%)		
INFLAMMATION, GRANULOMATOUS	1 (3%)		
INFLAMMATION, NECRO GRAN	1 (3%)		
NECROSIS, FOCAL	1 (3%)		
CYTOPLASMIC VACUOLIZATION	1 (3%)		
HEPATOCYTOMEGALY	2 (7%)		
HEMATOPOIESIS	1 (3%)		
*LIVER/CENTRILOBULAR	(30)	(33)	
NECROSIS, NOS		1 (3%)	
NECROSIS, COAGULATIVE	1 (3%)		
*PANCREAS	(30)	(33)	
ATROPHY, NOS		1 (3%)	
*COLON	(29)	(30)	
ULCER, NOS	1 (3%)		
*RECTUM	(30)	(33)	(34)
HEMATOMA, NOS		1 (3%)	
URINARY SYSTEM			
*KIDNEY	(30)	(33)	
HYDRONEPHROSIS	1 (3%)		
INFLAMMATION, INTERSTITIAL		1 (3%)	
GLOMERULONEPHRITIS, MEMBRANOUS		1 (3%)	
INFLAMMATION, CHRONIC	10 (33%)		
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C4. FEMALE RATS (TREATED GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
ENDOCRINE SYSTEM			
•ADRENAL	(30)	(33)	
ANGIECTASIS	11 (37%)	4 (12%)	
REPRODUCTIVE SYSTEM			
•MAMMARY GLAND	(30)	(33)	(34)
CYST, NOS	13 (43%)	3 (9%)	
INFLAMMATION, VESICULAR		1 (3%)	
•VAGINA	(30)	(33)	(34)
EPIDERMAL INCLUSION CYST	1 (3%)		
INFLAMMATION, CHRONIC SUPPURATIV	1 (3%)		
•UTERUS	(30)	(32)	
HEMORRHAGE	1 (3%)		
ANGIECTASIS	1 (3%)		
METAPLASIA, SQUAMOUS	1 (3%)		
•UTERUS/ENDOMETRIUM	(30)	(32)	
INFLAMMATION, SUPPURATIVE	4 (13%)	4 (13%)	
ULCER, CHRONIC	1 (3%)		
INFLAMMATION, CHRONIC SUPPURATIV	6 (20%)	2 (6%)	
HYPERPLASIA, CYSTIC	2 (7%)		
•OVARY	(30)	(32)	
CYST, NOS	14 (47%)		
INFLAMMATION, SUPPURATIVE		1 (3%)	
INFLAMMATION, CHRONIC SUPPURATIV	1 (3%)		
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
•EAR CANAL	(30)	(33)	(34)
KERATIN-PEARL FORMATION		1 (3%)	
MUSCULOSKELETAL SYSTEM			
NONE			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C4. FEMALE RATS (TREATED GROUPS): NONNEOPLASTIC LESIONS (CONTINUED)

	LOW DOSE	MID DOSE	HIGH DOSE
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED		1	
NECROPSY PERF/NO HISTO PERFORMED			34
AUTOLYSIS/NO NECROPSY	1	2	1
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

APPENDIX D

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MICE GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

TABLE D1.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MICE
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA**

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	15	15	@ 34	35
ANIMALS NECROPSIED	14	15	30	34
ANIMALS EXAMINED HISTOPATHOLOGICALLY	14	15	30	33
INTEGUMENTARY SYSTEM				
*SKIN	(14)	(15)	(30)	(34)
EPIDERMAL INCLUSION CYST			1 (3%)	
INFLAMMATION, CHRONIC SUPPURATIV	1 (7%)		1 (3%)	
*SUBCUT TISSUE	(14)	(15)	(30)	(34)
EDEMA, NOS				2 (6%)
INFLAMMATION, CHRONIC SUPPURATIV		1 (7%)		
METAPLASIA, SQUAMOUS			1 (3%)	
RESPIRATORY SYSTEM				
*LUNG	(14)	(15)	(30)	(33)
INFLAMMATION, INTERSTITIAL				1 (3%)
BRONCHOPNEUMONIA SUPPURATIVE		1 (7%)		
PERIARTERITIS			1 (3%)	
HEMATOPOIETIC SYSTEM				
*BONE MARROW	(13)	(14)	(29)	(32)
ATROPHY, NOS				4 (13%)
HYPERPLASIA, GRANULOCYTIC			1 (3%)	
*SPLEEN	(14)	(15)	(30)	(31)
HYPERPLASIA, GRANULOCYTIC				1 (3%)
HYPERPLASIA, LYMPHOID			2 (7%)	
HEMATOPOIESIS	1 (7%)	2 (13%)	18 (60%)	10 (32%)
*MESENTERIC L. NODE	(2)	(5)	(1)	(15)
CONGESTION, NOS		4 (80%)		
HYPERPLASIA, LYMPHOID	1 (50%)			

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

@NOTE: 35 ANIMALS WERE INITIALLY IN STUDY BUT ONE WAS FOUND TO BE A FEMALE IN A MALE GROUP AND WAS DELETED.

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
CIRCULATORY SYSTEM				
#MYOCARDIUM	(14)	(15)	(30)	(32)
INFLAMMATION, INTERSTITIAL			1 (3%)	
INFLAMMATION, CHRONIC			1 (3%)	
DIGESTIVE SYSTEM				
#LIVER	(14)	(15)	(30)	(33)
MINERALIZATION			1 (3%)	
CONGESTION, NOS		1 (7%)		
NECROSIS, NOS			1 (3%)	
NECROSIS, FOCAL				1 (3%)
HYPERPLASIA, NODULAR		3 (20%)		
#LIVER/CENTRIOLOBULAR	(14)	(15)	(30)	(33)
NECROSIS, NOS			1 (3%)	
#LIVER/PERIPORTAL	(14)	(15)	(30)	(33)
FIBROSIS			1 (3%)	
URINARY SYSTEM				
#KIDNEY	(14)	(15)	(30)	(33)
HYDRONEPHROSIS			1 (3%)	
INFLAMMATION, SUPPURATIVE				1 (3%)
PYELONEPHRITIS SUPPURATIVE	1 (7%)			
INFLAMMATION, CHRONIC			1 (3%)	
HYPERPLASIA, LYMPHOID			1 (3%)	
#URINARY BLADDER	(14)	(15)	(30)	(33)
INFLAMMATION, ACUTE/CHRONIC	1 (7%)			
INFLAMMATION, CHRONIC SUPPURATIVE	1 (7%)			
ENDOCRINE SYSTEM				
#ADRENAL	(14)	(15)	(30)	(33)
ANGIECTASIS			1 (3%)	
REPRODUCTIVE SYSTEM				
#PROSTATE	(14)	(15)	(30)	(33)
INFLAMMATION, SUPPURATIVE			2 (7%)	1 (3%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ABSCCESS, NOS			1 (3%)	
*TESTIS ATROPHY, NOS	(14)	(15)	(30)	(33) 1 (3%)
*TUNICA ALBUGINEA MINERALIZATION	(14)	(15)	(30) 1 (3%)	(33)
NERVOUS SYSTEM				
NONE				
SPECIAL SENSE ORGANS				
*EAE CANAL KERATIN-PEARL FORMATION	(14)	(15)	(30) 1 (3%)	(34)
MUSCULOSKELETAL SYSTEM				
NONE				
BODY CAVITIES				
NONE				
ALL OTHER SYSTEMS				
*MULTIPLE ORGANS HYPERPLASIA, LYMPHOID	(14)	(15)	(30) 1 (3%)	(34)
SPECIAL MORPHOLOGY SUMMARY				
NO LESION REPORTED	7	5	6	1
NECROPSY PERF/NO HISTO PERFORMED				1
AUTOLYSIS/NO NECROPSY	1		4	1
‡ NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY * NUMBER OF ANIMALS NECROPSIED				

TABLE D2.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MICE
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA**

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	15	15	35	35
ANIMALS MISSING			1	2
ANIMALS NECROPSIED	15	15	30	32
ANIMALS EXAMINED HISTOPATHOLOGICALLY	15	15	30	32
INTEGUMENTARY SYSTEM				
*SUBCUT TISSUE EDEMA, NOS	(15)	(15)	(30)	(32) 5 (16%)
RESPIRATORY SYSTEM				
#LUNG/BRONCHIOLE HYPERPLASIA, LYMPHOID	(15) 1 (7%)	(15) 1 (7%)	(29) 1 (3%)	(32)
#LUNG INFLAMMATION, INTERSTITIAL HYPERPLASIA, ALVEOLAR EPITHELIUM	(15)	(15)	(29) 2 (7%) 1 (3%)	(32)
HEMATOPOIETIC SYSTEM				
#BONE MARROW ATROPHY, NOS	(15) 1 (7%)	(15)	(29) 4 (14%)	(32)
#SPLEEN ANGIECTASIS HEMATOPOIESIS	(15) 3 (20%)	(15)	(30) 1 (3%) 5 (17%)	(32) 2 (6%)
#MESENTERIC L. NODE HYPERPLASIA, GRANULOCYTIC HYPERPLASIA, RETICULUM CELL HYPERPLASIA, LYMPHOID	(2) 1 (50%) 1 (50%)	(1)	(4) 1 (25%) 1 (25%)	(24)
#INGUINAL LYMPH NODE CONGESTION, NOS ATROPHY, NOS	(2) 1 (50%) 1 (50%)	(1)	(4)	(24)
CIRCULATORY SYSTEM				
#MYOCARDIUM INFLAMMATION, INTERSTITIAL	(15)	(15)	(29) 1 (3%)	(32)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
DIGESTIVE SYSTEM				
*LIVER	(15)	(15)	(30)	(32)
FIBROSIS, DIFFUSE			1 (3%)	
NECROSIS, FOCAL		1 (7%)		
INFARCT, NOS		1 (7%)		
HYPERPLASIA, NODULAR			1 (3%)	
ANGIECTASIS			1 (3%)	
HEMATOPOIESIS			1 (3%)	
*PANCREAS	(15)	(15)	(28)	(32)
ATROPHY, NOS	1 (7%)			
URINARY SYSTEM				
*KIDNEY	(15)	(15)	(30)	(32)
HYDRONEPHROSIS	1 (7%)			
INFLAMMATION, INTERSTITIAL	1 (7%)			
AMYLOIDOSIS			1 (3%)	
HYPERPLASIA, LYMPHOID	1 (7%)			
*URINARY BLADDER	(15)	(15)	(29)	(32)
INFLAMMATION, CHRONIC SUPPURATIV	1 (7%)			
ENDOCRINE SYSTEM				
NONE				
REPRODUCTIVE SYSTEM				
*UTERUS	(15)	(15)	(29)	(31)
EDEMA, NOS	1 (7%)			
*UTERUS/ENDOMETRIUM	(15)	(15)	(29)	(31)
INFLAMMATION, CHRONIC SUPPURATIV		1 (7%)		
HYPERPLASIA, CYSTIC	10 (67%)	10 (67%)	9 (31%)	
*OVARY	(15)	(15)	(29)	(31)
CYST, NOS			2 (7%)	1 (3%)
HEMORRHAGE			1 (3%)	
ATROPHY, NOS			1 (3%)	
NERVOUS SYSTEM				
NONE				

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	VEHICLE CONTROL	UNTREATED CONTROL	LOW DOSE	HIGH DOSE
SPECIAL SENSE ORGANS				
*EAR CANAL INFLAMMATION, CHRONIC SUPPURATIV	(15)	(15)	(30) 1 (3%)	(32)
MUSCULOSKELETAL SYSTEM				
NONE				
BODY CAVITIES				
*PERITONEUM INFLAMMATION, NOS	(15) 1 (7%)	(15)	(30)	(32)
ALL OTHER SYSTEMS				
*MULTIPLE ORGANS PERIARTERITIS HYPERPLASIA, LYMPHOID	(15)	(15) 1 (7%)	(30) 1 (3%)	(32)
SPECIAL MORPHOLOGY SUMMARY				
NO LESION REPORTED	4	3	6	
ANIMAL MISSING/NO NECROPSY			1	2
AUTO/NECROPSY/HISTO PERF			1	
AUTOLYSIS/NO NECROPSY			4	1
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY				
* NUMBER OF ANIMALS NECROPSIED				

APPENDIX E

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS IN RATS
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA

Table El. Analyses of the Incidence of Primary Tumors in Low-Dose
Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u> <u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>	<u>Low Dose</u>
Skin: Squamous-cell Carcinomab	0/30 (0)	0/10 (0)	5/37 (14)
P Values ^{c,d}			P = 0.045**
Relative Risk (Pooled Control) ^e Lower Limit Upper Limit			Infinite 1.044 Infinite
Relative Risk (Vehicle Control) ^e Lower Limit Upper Limit			Infinite 0.381 Infinite
Weeks to First Observed Tumor	--	--	53
Subcutaneous Tissue: Sarcoma, NOS ^b	0/30 (0)	0/10 (0)	5/37 (14)
P Values ^{c,d}			P = 0.045**
Relative Risk (Pooled Control) ^e Lower Limit Upper Limit			Infinite 1.044 Infinite
Relative Risk (Vehicle Control) ^e Lower Limit Upper Limit			Infinite 0.381 Infinite
Weeks to First Observed Tumor	--	--	61

Table El. Analyses of the Incidence of Primary Tumors in Low-Dose
Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Subcutaneous Tissue:			
Sarcoma or Fibrosarcoma ^b		1/30 (3)	0/10 (0)
P Values ^{c,d}			6/37 (16)
			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			4.865
Upper Limit			0.642
			216.767
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.484
			Infinite
Weeks to First Observed Tumor		--	61
Subcutaneous Tissue: Fibroma ^b		0/30 (0)	2/37 (5)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.244
			Infinite
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.088
			Infinite
Weeks to First Observed Tumor		--	74

Table E1. Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>	<u>Low Dose</u>
Hematopoietic System: Granulocytic Leukemia ^b	0/30 (0)	0/10 (0)	2/37 (5)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.244
			Infinite
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.088
			Infinite
Weeks to First Observed Tumor	--	--	55
Hematopoietic System: Lymphoma or Lymphocytic Leukemia ^b	0/30 (0)	0/10 (0)	4/37 (11)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.767
			Infinite
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.279
			Infinite
Weeks to First Observed Tumor	--	--	37

Table E1. Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
Topography: Morphology	Pooled Control	Low-Dose Vehicle Control	Low Dose
Hematopoietic System: Leukemia or Lymphoma ^b	0/30 (0)	0/10 (0)	6/37 (16)
P Values ^{c,d}			P = 0.023**
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			1.325
Relative Risk (Vehicle Control) ^e			Infinite
Lower Limit			Infinite
Upper Limit			0.484
Weeks to First Observed Tumor	--	--	37
Pituitary: Chromophobe Adenoma ^b	7/26 (27)	3/9 (33)	1/28 (4)
P Values ^{c,d}			P = 0.038*(N) P = 0.019**(N)
Relative Risk (Pooled Control) ^e			
Lower Limit			0.133
Upper Limit			0.003
Relative Risk (Vehicle Control) ^e			0.931
Lower Limit			0.107
Upper Limit			0.002
Weeks to First Observed Tumor	--	82	1.186
			78

Table E1. Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
Topography:	Morphology	Pooled Control	Low-Dose Vehicle Control
Ear Canal:	Squamous-cell Carcinoma ^b	0/30 (0)	0/10 (0)
P Values ^{c,d}			3/37 (8)
Relative Risk (Pooled Control) ^e			N.S.
Lower Limit			Infinitesimal
Upper Limit			0.497
Relative Risk (Vehicle Control) ^e			Infinitesimal
Lower Limit			Infinitesimal
Upper Limit			0.181
Weeks to First Observed Tumor		--	74
Skin and Ear Canal:			
Squamous-cell Carcinoma ^b		0/30 (0)	0/10 (0)
P Values ^{c,d}			7/37 (19)
Relative Risk (Pooled Control) ^e			p = 0.011**
Lower Limit			Infinitesimal
Upper Limit			1.608
Relative Risk (Vehicle Control) ^e			Infinitesimal
Lower Limit			Infinitesimal
Upper Limit			0.587
Weeks to First Observed Tumor		--	53

Table El. Analyses of the Incidence of Primary Tumors in Low-Dose
Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Mesentery:	Sarcoma, NOS ^b	0/30 (0)	0/10 (0)
P Values ^{c,d}			
Relative Risk (Pooled Control) ^e			
Lower Limit			
Upper Limit			
Relative Risk (Vehicle Control) ^e			
Lower Limit			
Upper Limit			
Weeks to First Observed Tumor		--	--
			66

^aTreated group received dose of 0.7 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent).

^cBeneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$ for either control group; otherwise, not significant (N.S.) is indicated.

^dA negative value (N) indicates a lower incidence in a treated group than in a control group.

^eThe 95% confidence interval of the relative risk between each treated group and the specified control group.

Table E2. Analyses of the Incidence of Primary Tumors in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>	<u>Mid Dose</u>
Skin: Squamous-cell Carcinoma ^b	0/30 (0)	0/10 (0)	3/30 (10)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.614
Relative Risk (Vehicle Control) ^e			Infinite
Lower Limit			0.223
Upper Limit			Infinite
Weeks to First Observed Tumor		--	44
Hematopoietic System: Lymphocytic Leukemia ^b	0/30 (0)	0/10 (0)	5/30 (17)
P Values ^{c,d}			P = 0.026**
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.291
Relative Risk (Vehicle Control) ^e			Infinite
Lower Limit			0.471
Upper Limit			Infinite
Weeks to First Observed Tumor		--	40

Table E2. Analyses of the Incidence of Primary Tumors in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
Topography:	Morphology	Pooled Control	Mid-Dose Vehicle Control
Hematopoietic System:	All Leukemia ^b	0/30 (0)	0/10 (0)
P Values ^{c,d}			6/30 (20)
			P = 0.012**
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinitive
Upper Limit			1.638
			Infinitive
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinitive
Upper Limit			0.598
			Infinitive
Weeks to First Observed Tumor		--	40
Pituitary:	Chromophobe Adenoma ^b	2/29 (7)	0/10 (0)
P Values ^{c,d}			1/24 (4)
			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			0.604
Upper Limit			0.011
			10.830
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinitive
Upper Limit			0.024
			Infinitive
Weeks to First Observed Tumor		--	54

Table E2. Analyses of the Incidence of Primary Tumors in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)		Mid-Dose Vehicle Control	Pooled Control	Mid-Dose
Topography: Morphology				Dose
Skin and Ear Canal:				
Squamous-cell Carcinoma ^b		0/30 (0)	0/10 (0)	3/30 (10)
P Values ^{c,d}				N.S.
Relative Risk (Pooled Control) ^e				
Lower Limit				Infinite
Upper Limit				0.614
				Infinite
Relative Risk (Vehicle Control) ^e				
Lower Limit				Infinite
Upper Limit				0.223
				Infinite
Weeks to First Observed Tumor		--		44

^aTreated group received dose of 1.4 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent).

^cBeneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$ for either control group; otherwise, not significant (N.S.) is indicated.

^dA negative value (N) indicates a lower incidence in a treated group than in a control group.

^eThe 95% confidence interval of the relative risk between each treated group and the specified control group.

Table E3. Analyses of the Incidence of Primary Tumors in Low-Dose
Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>	<u>Low Dose</u>
Subcutaneous Tissue: Sarcoma, NOS, or Fibrosarcoma ^b		0/30 (0)	0/10 (0)	2/30 (7)
P Values ^{c,d}				N.S.
Relative Risk (Pooled Control) ^e				Infinit
Lower Limit				0.301
Upper Limit				Infinit
Relative Risk (Vehicle Control) ^e				Infinit
Lower Limit				0.109
Upper Limit				Infinit
Weeks to First Observed Tumor		--	--	79
Hematopoietic System: Leukemia or Lymphoma ^b		1/30 (3)	0/10 (0)	2/30 (7)
P Values ^{c,d}				N.S.
Relative Risk (Pooled Control) ^e				2.000
Lower Limit				0.110
Upper Limit				113.910
Relative Risk (Vehicle Control) ^e				Infinit
Lower Limit				0.109
Upper Limit				Infinit
Weeks to First Observed Tumor		--	--	63

Table E3. Analyses of the Incidence of Primary Tumors in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Pituitary:	Chromophobe Adenoma ^b	14/28 (50)	1/8 (13)
P Values ^{c,d}			4/29 (14)
			P = 0.004**(N)
Relative Risk (Pooled Control) ^e			
Lower Limit			0.276
Upper Limit			0.078
			0.752
Relative Risk (Vehicle Control) ^e			1.103
Lower Limit			0.142
Upper Limit			52.317
Weeks to First Observed Tumor		--	82
			75
Ear Canal:	Squamous-cell Carcinoma ^b	0/30 (0)	0/10 (0)
P Values ^{c,d}			2/30 (7)
			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.301
			Infinite
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.109
			Infinite
Weeks to First Observed Tumor		--	--
			74

Table E3. Analyses of the Incidence of Primary Tumors in Low-Dose
Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Brain:	Olfactory Neuroblastoma ^b	0/30 (0)	0/10 (0)
P Values ^{c,d}			2/30 (7)
Relative Risk (Pooled Control) ^e			N.S.
Lower Limit			Infinite
Upper Limit			0.301
Relative Risk (Vehicle Control) ^e			Infinite
Lower Limit			Infinite
Upper Limit			0.109
Weeks to First Observed Tumor		---	59
Mammary Gland:	Adenocarcinoma, NOS ^b	1/30 (3)	0/10 (0)
P Values ^{c,d}			7/30 (23)
Relative Risk (Pooled Control) ^e			P = 0.026**
Lower Limit			7.000
Upper Limit			0.987
Relative Risk (Vehicle Control) ^e			302.176
Lower Limit			Infinite
Upper Limit			0.726
Weeks to First Observed Tumor		---	Infinite
			37

Table E3. Analyses of the Incidence of Primary Tumors in Low-Dose
Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
Topography: Morphology	Pooled Control	Low-Dose Vehicle Control	Low Dose
Mammary Gland: Fibroadenoma ^b	6/30 (20)	2/10 (20)	8/30 (27)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			1.333
Upper Limit			0.464
			4.085
Relative Risk (Vehicle Control) ^e			
Lower Limit			1.333
Upper Limit			0.348
			11.664
Weeks to First Observed Tumor	--	82	66
Uterus: Adenocarcinoma, NOS ^b	0/30 (0)	0/10 (0)	2/30 (7)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.301
			Infinite
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.109
			Infinite
Weeks to First Observed Tumor	--	--	70

Table E3. Analyses of the Incidence of Primary Tumors in Low-Dose
Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Uterus:	Sarcoma, NOS ^b	0/30 (0)	0/10 (0)
P Values ^{c,d}			2/30 (7)
			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			Infinite
Upper Limit			0.301
			Infinite
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.109
			Infinite
Weeks to First Observed Tumor		--	--
			82
Uterus:	Endometrial Stromal Polyp ^b	1/30 (3)	0/10 (0)
P Values ^{c,d}			4/30 (13)
			N.S.
Relative Risk (Pooled Control) ^e			
Lower Limit			4.000
Upper Limit			0.428
			189.625
Relative Risk (Vehicle Control) ^e			
Lower Limit			Infinite
Upper Limit			0.345
			Infinite
Weeks to First Observed Tumor		--	--
			63

Table E3. Analyses of the Incidence of Primary Tumors in Low-Dose
Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

^aTreated group received dose of 0.7 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent).

^cBeneath the incidence of tumors in a treated groups is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$ for either control group; otherwise, not significant (N.S.) is indicated.

^dA negative value (N) indicates a lower incidence in a treated group than in a control group.

^eThe 95% confidence interval of the relative risk between each treated group and the specified control group.

Table E4. Analyses of the Incidence of Primary Tumors in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>	<u>Mid Dose</u>
Skin: Squamous-cell Carcinoma ^b	0/28 (0)	0/10 (0)	3/33 (9)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			Infinitesimal
Lower Limit			0.522
Upper Limit			Infinitesimal
Relative Risk (Vehicle Control) ^e			Infinitesimal
Lower Limit			0.203
Upper Limit			Infinitesimal
Weeks to First Observed Tumor		--	64
Hematopoietic System: Leukemia or Lymphoma ^b	0/28 (0)	0/10 (0)	3/33 (9)
P Values ^{c,d}			N.S.
Relative Risk (Pooled Control) ^e			Infinitesimal
Lower Limit			0.522
Upper Limit			Infinitesimal
Relative Risk (Vehicle Control) ^e			Infinitesimal
Lower Limit			0.203
Upper Limit			Infinitesimal
Weeks to First Observed Tumor		--	25

Table E4. Analyses of the Incidence of Primary Tumors in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)		Mid-Dose	
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>
Pituitary:	Chromophobe Adenoma ^b	8/28 (29)	2/10 (20)
P Values ^{c,d}			0/32 (0)
			P = 0.001**(N)
Relative Risk (Pooled Control) ^e			0.000
Lower Limit			0.000
Upper Limit			0.374
Relative Risk (Vehicle Control) ^e			0.000
Lower Limit			0.000
Upper Limit			1.028
Weeks to First Observed Tumor			--
			87
Mammary Gland:	Adenoma, NOS ^b	0/28 (0)	0/10 (0)
P Values ^{c,d}			3/33 (9)
			N.S.
Relative Risk (Pooled Control) ^e			Infinite
Lower Limit			0.522
Upper Limit			Infinite
Relative Risk (Vehicle Control) ^e			Infinite
Lower Limit			0.203
Upper Limit			Infinite
Weeks to First Observed Tumor		--	43

Table E4. Analyses of the Incidence of Primary Tumors in Mid-Dose
Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

<u>Topography:</u> <u>Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>	<u>Mid Dose</u>
Mammary Gland: Adenocarcinoma, NOS ^b	1/28 (4)	1/10 (10)	8/33 (24)
P Values ^{c,d}			P = 0.025**
Relative Risk (Pooled Control) ^e			6.788
Lower Limit			1.004
Upper Limit			289.508
Relative Risk (Vehicle Control) ^e			2.424
Lower Limit			0.409
Upper Limit			103.495
Weeks to First Observed Tumor		87	40
Mammary Gland: Fibroadenoma ^b	10/28 (36)	5/10 (50)	4/33 (12)
P Values ^{c,d}			P = 0.020*(N) P = 0.030**(N)
Relative Risk (Pooled Control) ^e			0.339
Lower Limit			0.088
Upper Limit			1.036
Relative Risk (Vehicle Control) ^e			0.242
Lower Limit			0.071
Upper Limit			0.948
Weeks to First Observed Tumor		87	56

Table E4. Analyses of the Incidence of Primary Tumors in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)		Pooled Control	Mid-Dose Vehicle Control	Mid Dose
<u>Topography:</u> <u>Morphology</u>				
Uterus:	Adenocarcinoma, NOS ^b	0/28 (0)	0/10 (0)	7/32 (22)
P Values ^{c,d}				P = 0.009**
Relative Risk (Pooled Control) ^e				
	Lower Limit			Infinite
	Upper Limit			1.745
Relative Risk (Vehicle Control) ^e				
	Lower Limit			Infinite
	Upper Limit			0.680
Weeks to First Observed Tumor			--	50
Ear Canal: Squamous-cell Carcinoma ^b		0/28 (0)	0/10 (0)	5/33 (15)
P Values ^{c,d}				P = 0.040**
Relative Risk (Pooled Control) ^e				
	Lower Limit			Infinite
	Upper Limit			1.097
Relative Risk (Vehicle Control) ^e				
	Lower Limit			Infinite
	Upper Limit			0.427
Weeks to First Observed Tumor			--	71

Table E4. Analyses of the Incidence of Primary Tumors in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)		Mid-Dose Vehicle Control	Mid-Dose
Topography:	Morphology	Pooled Control	Dose
Skin and Ear Canal:			
	Squamous-cell Carcinoma ^b	0/28 (0)	8/33 (24)
P Values ^{c,d}			P = 0.005**
Relative Risk (Pooled Control) ^e			
	Lower Limit		Infinite
	Upper Limit		1.991
			Infinite
Relative Risk (Vehicle Control) ^e			
	Lower Limit		Infinite
	Upper Limit		0.776
			Infinite
Weeks to First Observed Tumor		--	64

^aTreated group received dose of 1.4 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent).

^cBeneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$ for either control group; otherwise, not significant (N.S.) is indicated.

^dA negative value (N) indicates a lower incidence in a treated group than in a control group.

^eThe 95% confidence interval of the relative risk between each treated group and the specified control group.

Table E5. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>	<u>Low Dose</u>
Skin:	Squamous-cell Carcinoma ^c	0/29 (0)	0/10 (0)	5/33 (15)
P Values ^{d,e}				P = 0.036**
Relative Risk (Pooled Control) ^f				
Lower Limit				Infinite
Upper Limit				1.134
				Infinite
Relative Risk (Vehicle Control) ^f				
Lower Limit				Infinite
Upper Limit				0.427
				Infinite
Weeks to First Observed Tumor			--	53
Subcutaneous Tissue:	Sarcoma, NOS ^c	0/29 (0)	0/10 (0)	5/33 (15)
P Values ^{d,e}				P = 0.036**
Relative Risk (Pooled Control) ^f				
Lower Limit				Infinite
Upper Limit				1.134
				Infinite
Relative Risk (Vehicle Control) ^f				
Lower Limit				Infinite
Upper Limit				0.427
				Infinite
Weeks to First Observed Tumor			--	61

Table E5. Time-adjusted Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
Topography: Morphology	Pooled Control	Low-Dose Vehicle Control	Low Dose
Subcutaneous Tissue: Sarcoma or Fibrosarcoma ^c	1/30 (3)	0/10 (0)	6/33 (18)
P Values ^{d,e}			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			5.455
Upper Limit			0.722
			241.878
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinte
Upper Limit			0.543
			Infinte
Weeks to First Observed Tumor		--	61
Subcutaneous Tissue: Fibroma ^c	0/29 (0)	0/10 (0)	2/33 (6)
P Values ^{d,e}			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinte
Upper Limit			0.265
			Infinte
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinte
Upper Limit			0.099
			Infinte
Weeks to First Observed Tumor		--	74

Table E5. Time-adjusted Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Hematopoietic System:			
Granulocytic Leukemia ^c			
		0/29 (0)	0/10 (0)
P Values ^{d,e}			
Relative Risk (Pooled Control) ^f			
Lower Limit			
Upper Limit			
Relative Risk (Vehicle Control) ^f			
Lower Limit			
Upper Limit			
Relative Risk (Vehicle Control) ^f			
Lower Limit			
Upper Limit			
Weeks to First Observed Tumor			
--			
55			
Hematopoietic System: Lymphoma or Lymphocytic Leukemia ^b			
		0/29 (0)	0/10 (0)
P Values ^{d,e}			
Relative Risk (Pooled Control) ^f			
Lower Limit			
Upper Limit			
Relative Risk (Vehicle Control) ^f			
Lower Limit			
Upper Limit			
Weeks to First Observed Tumor			
--			
37			

Table E5. Time-adjusted Analyses of the Incidence of Primary Tumors in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>	<u>Low Dose</u>
Hematopoietic System: Leukemia or Lymphoma ^b	0/29 (0)	0/10 (0)	6/34 (18)
P Values ^{d,e}			P = 0.020**
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinitive
Upper Limit			1.397
			Infinitive
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinitive
Upper Limit			0.527
			Infinitive
Weeks to First Observed Tumor	--	--	37
Pituitary: Chromophobe Adenoma ^c	7/25 (28)	3/9 (33)	1/27 (4)
P Values ^{d,e}			P = 0.041*(N) P = 0.019**(N)
Relative Risk (Pooled Control) ^f			
Lower Limit			0.132
Upper Limit			0.003
			0.925
Relative Risk (Vehicle Control) ^f			
Lower Limit			0.111
Upper Limit			0.003
			1.227
Weeks to First Observed Tumor		82	78

Table E5. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)		Low-Dose	
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low Dose</u>
Ear Canal:	Squamous-cell Carcinoma ^c	0/29 (0)	0/10 (0)
P Values ^{d,e}			3/33 (9)
			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinite
Upper Limit			0.540
			Infinite
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinite
Upper Limit			0.203
			Infinite
Weeks to First Observed Tumor		---	74
Skin or Ear Canal:			
Squamous-cell Carcinoma ^c		0/29 (0)	0/10 (0)
P Values ^{d,e}			7/33 (21)
			P = 0.009**
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinite
Upper Limit			1.748
			Infinite
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinite
Upper Limit			0.659
			Infinite
Weeks to First Observed Tumor		--	53

Table E5. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)		Low-Dose	
Topography:	Morphology	Pooled Control	Vehicle Control
Mesentery:	Sarcoma, NOS ^c	0/29 (0)	0/10 (0)
P Values ^{d,e}			2/33 (6)
			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinit
Upper Limit			0.265
			Infinit
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinit
Upper Limit			0.099
			Infinit
Weeks to First Observed Tumor		--	66

^aTreated group received dose of 0.7 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 37 weeks of the study.

^cNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 52 weeks of the study.

^dBeneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$; otherwise, not significant (N.S.) is indicated.

^eA negative value (N) indicates a lower incidence in a treated group than in a control group.

^fThe 95% confidence interval of the relative risk between each treated group and the control group.

Table E6. Time-adjusted Analyses of the Incidence of Primary Tumors
in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>	<u>Mid Dose</u>
Hematopoietic System: Lymphocytic Leukemia ^b	0/30 (0)	0/10 (0)	5/16 (31)
P Values ^{e,f}			P = 0.003**
Relative Risk (Pooled Control) ^g Lower Limit Upper Limit			Infinite 2.453 Infinite
Relative Risk (Vehicle Control) ^g Lower Limit Upper Limit			Infinite 0.892 Infinite
Weeks to First Observed Tumor		--	40
Hematopoietic System: All Leukemia ^b	0/30 (0)	0/10 (0)	6/16 (38)
P Values ^{e,f}			P = 0.035* P = 0.001**
Relative Risk (Pooled Control) ^g Lower Limit Upper Limit			Infinite 3.123 Infinite
Relative Risk (Vehicle Control) ^g Lower Limit Upper Limit			Infinite 1.135 Infinite
Weeks to First Observed Tumor		--	40

Table E6. Time-adjusted Analyses of the Incidence of Primary Tumors in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>
Pituitary:	Chromophobe Adenoma ^d	2/29 (7)	0/10 (0)
P Values ^{e,f}			1/6 (17)
			N.S.
Relative Risk (Pooled Control) ^g			
Lower Limit			2.417
Upper Limit			0.042
			34.031
Relative Risk (Vehicle Control) ^g			
Lower Limit			Infinite
Upper Limit			0.096
			Infinite
Weeks to First Observed Tumor		--	54
Skin or Ear Canal:			
Squamous-cell Carcinoma ^c		0/30 (0)	0/10 (0)
P Values ^{e,f}			3/13 (23)
			P = 0.023**
Relative Risk (Pooled Control) ^g			
Lower Limit			Infinite
Upper Limit			1.435
			Infinite
Relative Risk (Vehicle Control) ^g			
Lower Limit			Infinite
Upper Limit			0.521
			Infinite
Weeks to First Observed Tumor		--	44

Table E6. Time-adjusted Analyses of the Incidence of Primary Tumors
in Mid-Dose Male Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

^aTreated group received dose of 1.4 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 40 weeks of the study.

^cNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 44 weeks of the study.

^dNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 52 weeks of the study.

^eBeneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$; otherwise, not significant (N.S.) is indicated.

^fA negative value (N) indicates a lower incidence in a treated group than in a control group.

^gThe 95% confidence interval of the relative risk between each treated group and the control group.

Table E7. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>	<u>Low Dose</u>
Subcutaneous Tissue: Sarcoma, NOS, or Fibrosarcoma ^c	0/28 (0)	0/9 (0)	2/29 (7)
P Values ^{d,e}			N.S.
Relative Risk (Pooled Control) ^f			Infinit
Lower Limit			0.292
Upper Limit			Infinit
Relative Risk (Vehicle Control) ^f			Infinit
Lower Limit			0.103
Upper Limit			Infinit
Weeks to First Observed Tumor	--	--	79
Hematopoietic System: Leukemia or Lymphoma ^c	1/28 (4)	0/9 (0)	2/29 (7)
P Values ^{d,e}			N.S.
Relative Risk (Pooled Control) ^f			1.931
Lower Limit			0.107
Upper Limit			109.859
Relative Risk (Vehicle Control) ^f			Infinit
Lower Limit			0.103
Upper Limit			Infinit
Weeks to First Observed Tumor	--	--	63

Table E7. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Pituitary:	Chromophobe Adenoma ^c	14/28 (50)	1/8 (13)
P Values ^{d,e}			4/28 (14)
			P = 0.005**(N)
Relative Risk (Pooled Control) ^f			
Lower Limit			0.286
Upper Limit			0.087
			0.777
Relative Risk (Vehicle Control) ^f			
Lower Limit			1.143
Upper Limit			0.147
			54.086
Weeks to First Observed Tumor		82	75
Ear Canal:	Squamous-cell Carcinoma ^c	0/28 (0)	2/29 (7)
P Values ^{d,e}			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinite
Upper Limit			0.292
			Infinite
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinite
Upper Limit			0.103
			Infinite
Weeks to First Observed Tumor		---	74

Table E7. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
Topography:	Morphology	Pooled Control	Low-Dose Vehicle Control
Brain:	Olfactory Neuroblastoma ^c	0/28 (0)	0/9 (0)
P Values ^{d,e}			2/29 (7)
			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinitive
Upper Limit			0.292
			Infinitive
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinitive
Upper Limit			0.103
			Infinitive
Weeks to First Observed Tumor		--	59
Mammary Gland:	Adenocarcinoma, NOS ^b	1/28 (4)	0/9 (0)
P Values ^{d,e}			7/30 (23)
			P = 0.033**
Relative Risk (Pooled Control) ^f			
Lower Limit			6.533
Upper Limit			0.926
			282.066
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinitive
Upper Limit			0.663
			Infinitive
Weeks to First Observed Tumor		--	37

Table E7. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Mammary Gland:	Fibroadenoma ^c	6/28 (21)	2/9 (22)
P Values ^{d,e}			8/29 (28)
			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			1.287
Upper Limit			0.452
			3.917
Relative Risk (Vehicle Control) ^f			
Lower Limit			1.241
Upper Limit			0.335
			10.781
Weeks to First Observed Tumor		82	66
Uterus: Adenocarcinoma, NOS ^c		0/28 (0)	2/29 (7)
P Values ^{d,e}			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinite
Upper Limit			0.292
			Infinite
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinite
Upper Limit			0.103
			Infinite
Weeks to First Observed Tumor		--	70

Table E7. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Low-Dose Vehicle Control</u>
Uterus:	Sarcoma, NOS ^c	0/28 (0)	0/9 (0)
P Values ^{d,e}			2/29 (7)
			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			Infinitesimal
Upper Limit			0.292
			Infinitesimal
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinitesimal
Upper Limit			0.103
			Infinitesimal
Weeks to First Observed Tumor		--	82
Uterus:	Endometrial Stromal Polyp ^c	1/28 (4)	0/9 (0)
P Values ^{d,e}			4/29 (14)
			N.S.
Relative Risk (Pooled Control) ^f			
Lower Limit			3.862
Upper Limit			0.416
			182.833
Relative Risk (Vehicle Control) ^f			
Lower Limit			Infinitesimal
Upper Limit			0.326
			Infinitesimal
Weeks to First Observed Tumor		--	63

Table E7. Time-adjusted Analyses of the Incidence of Primary Tumors
in Low-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

^aTreated group received dose of 0.7 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 37 weeks of the study.

^cNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 52 weeks of the study.

^dBeneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$; otherwise, not significant (N.S.) is indicated.

^eA negative value (N) indicates a lower incidence in a treated group than in a control group.

^fThe 95% confidence interval of the relative risk between each treated group and the control group.

Table E8. Time-adjusted Analyses of the Incidence of Primary Tumors
in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>	<u>Mid Dose</u>
Skin:	Squamous-cell Carcinoma ^f	0/28 (0)	0/10 (0)	3/21 (14)
P Values ^{g,h}				N.S.
	Relative Risk (Pooled Control) ⁱ			Infinit
	Lower Limit			0.824
	Upper Limit			Infinit
	Relative Risk (Vehicle Control) ⁱ			Infinit
	Lower Limit			0.320
	Upper Limit			Infinit
Weeks to First Observed Tumor			---	64
Hematopoietic System: Lymphocytic				
	Leukemia or Lymphoma ^b	0/28 (0)	0/10 (0)	3/22 (14)
P Values ^{g,h}				N.S.
	Relative Risk (Pooled Control) ⁱ			Infinit
	Lower Limit			0.786
	Upper Limit			Infinit
	Relative Risk (Vehicle Control) ⁱ			Infinit
	Lower Limit			0.305
	Upper Limit			Infinit
Weeks to First Observed Tumor			---	25

Table E8. Time-adjusted Analyses of the Incidence of Primary Tumors
in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
Topography: Morphology	Pooled Control	Mid-Dose Vehicle Control	Mid Dose
Pituitary: Chromophobe Adenoma ^f	8/28 (29)	2/10 (20)	0/20 (0)
P Values ^{g,h}			P = 0.008**(N)
Relative Risk (Pooled Control) ⁱ			
Lower Limit			0.000
Upper Limit			0.000
			0.583
Relative Risk (Vehicle Control) ⁱ			0.000
Lower Limit			0.000
Upper Limit			1.609
Weeks to First Observed Tumor		87	--
Mammary Gland: Adenoma ^d	0/28 (0)	0/10 (0)	3/22 (14)
P Values ^{g,h}			N.S.
Relative Risk (Pooled Control) ⁱ			Infinite
Lower Limit			0.786
Upper Limit			Infinite
Relative Risk (Vehicle Control) ⁱ			Infinite
Lower Limit			0.305
Upper Limit			Infinite
Weeks to First Observed Tumor		--	43

Table E8. Time-adjusted Analyses of the Incidence of Primary Tumors
in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>
Mammary Gland:	Adenocarcinoma, NOS ^c	1/28 (4)	1/10 (10)
P Values ^{g,h}			8/24 (33)
			P = 0.006**
Relative Risk (Pooled Control) ⁱ			
Lower Limit			9.333
Upper Limit			1.398
			388.857
Relative Risk (Vehicle Control) ⁱ			
Lower Limit			3.333
Upper Limit			0.567
			139.059
Weeks to First Observed Tumor			87
			40
Mammary Gland:	Fibroadenoma ^f	10/28 (36)	5/10 (50)
P Values ^{g,h}			4/21 (19)
			N.S.
Relative Risk (Pooled Control) ⁱ			
Lower Limit			0.533
Upper Limit			0.141
			1.555
Relative Risk (Vehicle Control) ⁱ			
Lower Limit			0.381
Upper Limit			0.113
			1.434
Weeks to First Observed Tumor			87
			56

Table E8. Time-adjusted Analyses of the Incidence of Primary Tumors
in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
Topography:	Morphology	Pooled Control	Mid-Dose Vehicle Control	Mid Dose
Uterus:	Adenocarcinoma, NOSE	0/28 (0)	0/10 (0)	7/21 (33)
P Values ^{g,h}				P = 0.044* P = 0.001**
Relative Risk (Pooled Control) ⁱ				
Lower Limit				Infinite
Upper Limit				2.68 ⁴ Infinite
Relative Risk (Vehicle Control) ⁱ				
Lower Limit				Infinite
Upper Limit				1.04 ⁴ Infinite
Weeks to First Observed Tumor			--	50
Ear Canal:	Squamous-cell Carcinoma ^f	0/28 (0)	0/10 (0)	5/21 (24)
P Values ^{g,h}				P = 0.011**
Relative Risk (Pooled Control) ⁱ				
Lower Limit				Infinite
Upper Limit				1.73 ⁷ Infinite
Relative Risk (Vehicle Control) ⁱ				
Lower Limit				Infinite
Upper Limit				0.67 ⁶ Infinite
Weeks to First Observed Tumor			--	71

Table E8. Time-adjusted Analyses of the Incidence of Primary Tumors in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)			
<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Mid-Dose Vehicle Control</u>	<u>Mid Dose</u>
Skin and Ear Canal: Squamous-cell Carcinoma ^f	0/28 (0)	0/10 (0)	8/21 (38)
P Values ^{g,h}			P = 0.026* P < 0.001**
Relative Risk (Pooled Control) ⁱ			
Lower Limit			Infinite
Upper Limit			3.164
Relative Risk (Vehicle Control) ⁱ			Infinite
Lower Limit			Infinite
Upper Limit			1.230
Weeks to First Observed Tumor		--	64

^aTreated group received dose of 1.4 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 25 weeks of the study.

^cNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 40 weeks of the study.

^dNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 43 weeks of the study.

Table E8. Time-adjusted Analyses of the Incidence of Primary Tumors
in Mid-Dose Female Rats Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

eNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 50 weeks of the study.

fNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 52 weeks of the study.

gBeneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$; otherwise, not significant (N.S.) is indicated.

hA negative value (N) indicates a lower incidence in a treated group than in a control group.

iThe 95% confidence interval of the relative risk between each treated group and the control group.

APPENDIX F

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS IN MICE
GIVEN INTRAPERITONEAL INJECTIONS OF THIO-TEPA

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Skin:	Squamous-cell Carcinoma ^b	0/28 (0)	0/14 (0)	7/30 (23)	0/34 (0)
P Values ^{c,d}		N.S.	N.S.	P = 0.007**	N.S.
Departure from Linear Trend ^e		P < 0.001	P < 0.001		
Relative Risk (Pooled Control) ^f					
Lower Limit				Infinite	--
Upper Limit				1.863	--
				Infinite	--
Relative Risk (Vehicle Control) ^f					
Lower Limit				Infinite	--
Upper Limit				0.978	--
				Infinite	--
Weeks to First Observed Tumor		--	--	67	--

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u> <u>High Dose</u>
Lung:	Alveolar/Bronchiolar Adenoma ^b	1/28 (4)	1/14 (7)	5/30 (17) 0/33 (0)
P Values ^{c,d}		N.S.	N.S.	N.S.
Departure from Linear Trend ^e		P = 0.007	P = 0.043	
Relative Risk (Pooled Control) ^f				
Lower Limit				0.000
Upper Limit				0.000 15.686
Relative Risk (Vehicle Control) ^f				
Lower Limit				0.000
Upper Limit				0.000 7.858
Weeks to First Observed Tumor		--	87	83 --

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)					
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
	Hematopoietic System: Lymphoma or Lymphocytic Leukemia ^b	1/28 (4)	1/14 (7)	2/30 (7)	26/34 (76)
P Values ^{c,d}		P < 0.001	P < 0.001	N.S.	P < 0.001* P < 0.001**
Departure from Linear Trend ^e		P = 0.001	P = 0.003		
Relative Risk (Pooled Control) ^f					
Lower Limit				1.867	21.412
Upper Limit				0.103 106.333	4.109 770.826
Relative Risk (Vehicle Control) ^f					
Lower Limit				0.933	10.706
Upper Limit				0.055 53.204	2.187 385.659
Weeks to First Observed Tumor		--	78	61	26

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Hematopoietic System: Leukemia or Lymphoma ^b	1/28 (4)	1/14 (7)	3/30 (10)	26/34 (76)
P Values ^{c,d}	P < 0.001	P < 0.001	N.S.	P < 0.001* P < 0.001**
Departure from Linear Trend ^e	P = 0.003	P = 0.006		
Relative Risk (Pooled Control) ^f				
Lower Limit			2.800	21.412
Upper Limit			0.242 141.729	4.109 770.721
Relative Risk (Vehicle Control) ^f				
Lower Limit			1.400	10.706
Upper Limit			0.129 70.914	2.187 385.659
Weeks to First Observed Tumor	--	78	61	26

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
Topography: Morphology	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Liver: Hepatocellular Adenoma or Carcinoma ^b	1/28 (4)	1/14 (7)	4/30 (13)	0/33 (0)
P Values ^{c,d}	N.S.	N.S.	N.S.	N.S.
Departure from Linear Trend ^e	P = 0.023			
Relative Risk (Pooled Control) ^f				
Lower Limit			3.733	0.000
Upper Limit			0.401	0.000
			177.009	15.686
Relative Risk (Vehicle Control) ^f				
Lower Limit			1.867	0.000
Upper Limit			0.215	0.000
			88.558	7.858
Weeks to First Observed Tumor	--	87	67	--

Table Fl. Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u> <u>High Dose</u>
Ear Canal:	Squamous-cell			
	Carcinoma ^b	0/28 (0)	0/14 (0)	2/30 (7) 0/34 (0)
P Values ^{c,d}		N.S.	N.S.	N.S.
Departure from Linear Trend ^e		P = 0.040		
Relative Risk (Pooled Control) ^f				
Lower Limit				Infinitive
Upper Limit				0.282
				Infinitive
Relative Risk (Vehicle Control) ^f				
Lower Limit				Infinitive
Upper Limit				0.147
				Infinitive
Weeks to First Observed Tumor		---	---	83

Table Fl. Analyses of the Incidence of Primary Tumors in
Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u> <u>High Dose</u>
Preputial Gland:	Squamous-cell			
Carcinoma ^b		0/28 (0)	0/14 (0)	6/30 (20) 1/34 (3)
P Values ^{c,d}		N.S.	N.S.	P = 0.014** N.S.
Departure from Linear Trend ^e		P = 0.002	P = 0.007	
Relative Risk (Pooled Control) ^f				
Lower Limit				Infinite
Upper Limit				0.045
				Infinite
Relative Risk (Vehicle Control) ^f				
Lower Limit				Infinite
Upper Limit				0.023
				Infinite
Weeks to First Observed Tumor		--	--	72 56

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)		Pooled Control		Vehicle Control		Low Dose		High Dose	
Topography:	Morphology								
All Sites:	Squamous-cell Carcinoma ^b	0/28 (0)		0/14 (0)		14/30 (47)		1/34 (3)	
P Values ^{c,d}		N.S.		N.S.		P < 0.001** P = 0.001*		N.S.	
Departure from Linear Trend ^e		P < 0.001		P < 0.001					
Relative Risk (Pooled Control) ^f									
Lower Limit						Infinite		Infinite	
Upper Limit						4.211		0.045	
Relative Risk (Vehicle Control) ^f						Infinite		Infinite	
Lower Limit						Infinite		Infinite	
Upper Limit						2.209		0.023	
Weeks to First Observed Tumor		--		--		67		56	

Table Fl. Analyses of the Incidence of Primary Tumors in
Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

^aTreated groups received doses of 1.15 or 2.3 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent).

^cBeneath the incidence of tumors in a control group is the probability level for the Cochran-Armitage test when $P < 0.05$; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$ for either control group; otherwise, not significant (N.S.) is indicated.

^dA negative trend (N) indicates a lower incidence in a treated group than in a control group.

^eThe probability level for departure from linear trend is given when $P < 0.05$ for any comparison.

^fThe 95% confidence interval of the relative risk between each treated group and the specified control group.

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Lung:	Alveolar/Bronchiolar Adenoma ^b	0/30 (0)	0/15 (0)	5/29 (17)	0/32 (0)
P Values ^{c,d}		N.S.	N.S.	P = 0.024**	N.S.
Departure from Linear Trend ^e		P = 0.001	P = 0.005		
Relative Risk (Pooled Control) ^f					
Lower Limit				Infinite	--
Upper Limit				1.336	--
				Infinite	--
Relative Risk (Vehicle Control) ^f					
Lower Limit				Infinite	--
Upper Limit				0.698	--
				Infinite	--
Weeks to First Observed Tumor		--	--	86	--

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)					
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Hematopoietic System: Lymphocytic Leukemia or Lymphoma ^b		1/30 (3)	0/15 (0)	5/30 (17)	32/32 (100)
P Values ^{c,d}		P < 0.001	P < 0.001	N.S.	P < 0.001* P < 0.001**
Departure from Linear Trend ^e		P = 0.001	P = 0.038		
Relative Risk (Pooled Control) ^f					
Lower Limit				5.000	30.030
Upper Limit				0.609 227.307	8.208 Infinite
Relative Risk (Vehicle Control) ^f					
Lower Limit				Infinite	Infinite
Upper Limit				0.675 Infinite	6.468 Infinite
Weeks to First Observed Tumor		--	--	31	26

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)					
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Liver: Hepatocellular Adenoma ^b		0/30 (0)	0/15 (0)	2/30 (7)	0/32 (0)
P Values ^{c,d}		N.S.	N.S.	N.S.	N.S.
Departure from Linear Trend ^e		P = 0.040			
Relative Risk (Pooled Control) ^f					
Lower Limit				Infinite	--
Upper Limit				0.301	--
				Infinite	--
Relative Risk (Vehicle Control) ^f					
Lower Limit				Infinite	--
Upper Limit				0.157	--
				Infinite	--
Weeks to First Observed Tumor		--	--	86	--
Ovary: Carcinoma, NOS ^b		0/29 (0)	0/15 (0)	3/29 (10)	0/31 (0)
P Values ^{c,d}		N.S.	N.S.	N.S.	N.S.
Departure from Linear Trend ^e		P = 0.011	P = 0.030		
Relative Risk (Pooled Control) ^f					
Lower Limit				Infinite	--
Upper Limit				0.615	--
				Infinite	--
Relative Risk (Vehicle Control) ^f					
Lower Limit				Infinite	--
Upper Limit				0.332	--
				Infinite	--
Weeks to First Observed Tumor		--	--	78	--

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

^aTreated groups received doses of 1.15 or 2.3 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent).

^cBeneath the incidence of tumors in a control group is the probability level for the Cochran-Armitage test when $P < 0.05$; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$ for either control group; otherwise, not significant (N.S.) is indicated.

^dA negative trend (N) indicates a lower incidence in a treated group than in a control group.

^eThe probability level for departure from linear trend is given when $P < 0.05$ for any comparison.

^fThe 95% confidence interval of the relative risk between each treated group and the specified control group.

Table F3. Time-adjusted Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u> Morphology	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Skin: Squamous-cell Carcinoma ^d	0/18 (0)	0/8 (0)	7/24 (29)	0/2 (0)
P Values ^{e,f}	N.S.	N.S.	P = 0.013**	N.S.
Relative Risk (Pooled Control) ^h				
Lower Limit			Infinite	--
Upper Limit			1.544	--
			Infinite	--
Relative Risk (Matched Control) ^h				
Lower Limit			Infinite	--
Upper Limit			0.754	--
			Infinite	--
Weeks to First Observed Tumor		--	67	--
Lung: Alveolar/Bronchiolar Adenoma ^d	1/18 (6)	1/8 (13)	5/24 (21)	0/2 (0)
P Values ^{e,f}	N.S.	N.S.	N.S.	N.S.
Relative Risk (Pooled Control) ^h				
Lower Limit			3.750	0.000
Upper Limit			0.479	0.000
			168.323	93.466
Relative Risk (Matched Control) ^h				
Lower Limit			1.667	0.000
Upper Limit			0.246	0.000
			74.916	41.780
Weeks to First Observed Tumor		87	83	--

Table F3. Time-adjusted Analyses of the Incidence of Primary Tumors
in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
<u>Topography: Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Hematopoietic System: Lymphoma or Lymphocytic Leukemia ^b	1/18 (6)	1/8 (13)	2/24 (8)	26/28 (93)
P Values ^{e,f}	P < 0.001	P < 0.001	N.S.	P < 0.001* P < 0.001**
Departure from Linear Trend ^g	P < 0.001	P = 0.002		
Relative Risk (Pooled Control) ^h				
Lower Limit			1.500	16.714
Upper Limit			0.086 84.637	3.601 358.473
Relative Risk (Matched Control) ^h				
Lower Limit			0.667	7.429
Upper Limit			0.043 37.663	1.771 159.766
Weeks to First Observed Tumor		78	61	26

Table F3. Time-adjusted Analyses of the Incidence of Primary Tumors
in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)				
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>
				<u>High Dose</u>
Hematopoietic System:	Leukemia	1/18 (6)	1/8 (13)	3/24 (13)
or Lymphoma ^b				26/28 (93)
P Values ^{e,f}		P < 0.001	P < 0.001	P < 0.001*
				P < 0.001**
Departure from Linear Trend ^g		P = 0.003	P = 0.004	
Relative Risk (Pooled Control) ^h				
Lower Limit				2.250
Upper Limit				0.202
				112.716
Relative Risk (Matched Control) ^h				
Lower Limit				1.000
Upper Limit				0.103
				50.152
Weeks to First Observed Tumor			78	61
				26

Table 1 Time-adjusted Analyses of the Incidence of Primary Tumors in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)					
Topography:	Morphology	Pooled Control	Vehicle Control	Low Dose	High Dose
Liver: Hepatocellular Adenoma or Carcinoma ^d		1/18 (6)	1/8 (13)	4/24 (17)	0/2 (0)
P Values ^{e,f}		N.S.	N.S.	N.S.	N.S.
Relative Risk (Pooled Control) ^h					
Lower Limit				3.000	0.000
Upper Limit				0.336	0.000
				140.603	93.466
Relative Risk (Matched Control) ^h					
Lower Limit				1.333	0.000
Upper Limit				0.173	0.000
				62.559	41.780
Weeks to First Observed Tumor			87	67	--
Ear Canal: Squamous-cell Carcinoma ^d		0/18 (0)	0/8 (0)	2/24 (8)	0/2 (0)
P Values ^{e,f}		N.S.	N.S.	N.S.	N.S.
Relative Risk (Pooled Control) ^h					
Lower Limit				Infinite	--
Upper Limit				0.232	--
				Infinite	--
Relative Risk (Matched Control) ^h					
Lower Limit				Infinite	--
Upper Limit				0.113	--
				Infinite	--
Weeks to First Observed Tumor			--	83	--

Table F3. Time-adjusted Analyses of the Incidence of Primary Tumors
in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)						
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>	
Preputial Gland:	Squamous-cell					
	Carcinoma ^d	0/18 (0)	0/8 (0)	6/24 (25)	1/2 (50)	
P Values ^{e,f}		P = 0.014	N.S.	P = 0.026**	N.S.	
Relative Risk (Pooled Control) ^h						
	Lower Limit			Infinite	Infinite	
	Upper Limit			1.271	1.487	
				Infinite	Infinite	
Relative Risk (Matched Control) ^h						
	Lower Limit			Infinite	Infinite	
	Upper Limit			0.620	0.232	
				Infinite	Infinite	
Weeks to First Observed Tumor			--	72	56	
All Sites: Squamous-cell						
	Carcinoma ^d	0/18 (0)	0/8 (0)	14/24 (58)	1/2 (50)	
P Values ^{e,f}		P = 0.001	P = 0.018	P = 0.004*	N.S.	
				P < 0.001**		
Relative Risk (Pooled Control) ^h						
	Lower Limit			Infinite	Infinite	
	Upper Limit			3.513	0.487	
				Infinite	Infinite	
Relative Risk (Matched Control) ^h						
	Lower Limit			Infinite	Infinite	
	Upper Limit			1.709	0.232	
				Infinite	Infinite	
Weeks to First Observed Tumor			--	67	56	

Table F3. Time-adjusted Analyses of the Incidence of Primary Tumors
in Male Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)

^aTreated groups received doses of 1.15 or 2.3 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 26 weeks of the study.

^cNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 32 weeks of the study.

^dNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 52 weeks of the study.

^eBeneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when $P < 0.05$; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$; otherwise, not significant (N.S.) is indicated.

^fA negative trend (N) indicates a lower incidence in a treated group than in a control group.

^gThe probability level for departure from linear trend is given when $P < 0.05$ for any comparison.

^hThe 95% confidence interval of the relative risk between each treated group and the control group.

Table F4. Time-adjusted Analyses of the Incidence of Primary Tumors in Female Mice Given Intraperitoneal Injections of Thio-TEPA^a

<u>Topography:</u> <u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Lung: Alveolar/Bronchiolar Adenoma ^d	0/29 (0)	0/14 (0)	5/24 (21)	0/0 (-)
P Values ^{e,f}	P = 0.034	N.S.	P = 0.015**	N.S.
Relative Risk (Pooled Control) ^h				
Lower Limit			Infinitesimal	--
Upper Limit			1.567	--
			Infinitesimal	--
Relative Risk (Matched Control) ^h				
Lower Limit			Infinitesimal	--
Upper Limit			0.795	--
			Infinitesimal	--
Weeks to First Observed Tumor		--	86	--
Hematopoietic System: Lymphoma or Lymphocytic Leukemia ^b	0/29 (0)	0/14 (0)	5/26 (19)	32/32 (100)
P Values ^{e,f}	P < 0.001	P < 0.001	P = 0.019**	P < 0.001* P < 0.001**
Departure from Linear Trends	P = 0.006	P = 0.011		
Relative Risk (Pooled Control) ^h				
Lower Limit			Infinitesimal	Infinitesimal
Upper Limit			1.445	12.195
			Infinitesimal	Infinitesimal
Relative Risk (Matched Control) ^h				
Lower Limit			Infinitesimal	Infinitesimal
Upper Limit			0.733	6.060
			Infinitesimal	Infinitesimal
Weeks to First Observed Tumor		--	31	26

Table F4. Time-adjusted Analyses of the Incidence of Primary Tumors
in Female Mice Given Intraperitoneal Injections of Thio-TEPA^a

(continued)						
<u>Topography:</u>	<u>Morphology</u>	<u>Pooled Control</u>	<u>Vehicle Control</u>	<u>Low Dose</u>	<u>High Dose</u>	
Liver:	Hepatocellular Adenoma ^d	0/29 (0)	0/14 (0)	2/25 (8)	0/0 (-)	
P Values ^{e,f}		N.S.	N.S.	N.S.	N.S.	
Relative Risk (Pooled Control) ^h						
	Lower Limit			Infinite	--	
	Upper Limit			0.350	--	
				Infinite	--	
Relative Risk (Matched Control) ^h						
	Lower Limit			Infinite	--	
	Upper Limit			0.177	--	
				Infinite	--	
Weeks to First Observed Tumor			--	86	--	
Ovary:	Carcinoma, NOS ^d	0/28 (0)	0/14 (0)	3/24 (13)	0/0 (-)	
P Values ^{e,f}		N.S.	N.S.	N.S.	N.S.	
Relative Risk (Pooled Control) ^h						
	Lower Limit			Infinite	--	
	Upper Limit			0.720	--	
				Infinite	--	
Relative Risk (Matched Control) ^h						
	Lower Limit			Infinite	--	
	Upper Limit			0.377	--	
				Infinite	--	
Weeks to First Observed Tumor			--	78	--	

**Table F4. Time-adjusted Analyses of the Incidence of Primary Tumors
in Female Mice Given Intraperitoneal Injections of Thio-TEPA^a**

(continued)

^aTreated groups received doses of 1.15 or 2.3 mg/kg.

^bNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 26 weeks of the study.

^cNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 31 weeks of the study.

^dNumber of tumor-bearing animals/number of animals examined at site (percent) which survived at least 52 weeks of the study.

^eBeneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when $P < 0.05$; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a treated group is the probability level for the Fisher exact test for the comparison of that treated group with the vehicle-control group (*) or with the pooled-control group (**) when $P < 0.05$; otherwise, not significant (N.S.) is indicated.

^fA negative trend (N) indicates a lower incidence in a treated group than in a control group.

^gThe probability level for departure from linear trend is given when $P < 0.05$ for any comparison.

^hThe 95% confidence interval of the relative risk between each treated group and the control group.

Review of the Bioassay of Thio-TEPA* for Carcinogenicity
by the Data Evaluation/Risk Assessment Subgroup
of the Clearinghouse on Environmental Carcinogens

April 26, 1978

The Clearinghouse on Environmental Carcinogens was established in May, 1976, in compliance with DHEW Committee Regulations and the Provisions of the Federal Advisory Committee Act. The purpose of the Clearinghouse is to advise the Director of the National Cancer Institute (NCI) on its bioassay program to identify and to evaluate chemical carcinogens in the environment to which humans may be exposed. The members of the Clearinghouse have been drawn from academia, industry, organized labor, public interest groups, State health officials, and quasi-public health and research organizations. Members have been selected on the basis of their experience in carcinogenesis or related fields and, collectively, provide expertise in chemistry, biochemistry, biostatistics, toxicology, pathology, and epidemiology. Representatives of various Governmental agencies participate as ad hoc members. The Data Evaluation/Risk Assessment Subgroup of the Clearinghouse is charged with the responsibility of providing a peer review of reports prepared on NCI-sponsored bioassays of chemicals studied for carcinogenicity. It is in this context that the below critique is given on the bioassay of Thio-TEPA for carcinogenicity.

The primary reviewer said that, under the conditions of test, Thio-TEPA induced squamous-cell carcinomas of the skin and ear canal in both sexes of treated rats and hematopoietic neoplasms in the males. Lymphomas or lymphocytic leukemia were induced in both sexes of mice and squamous-cell carcinomas in the skin and associated glands of males. After a brief description of the experimental design and conditions of test, he noted the following points in his critique: the poor survival among the high dose treated animals resulting from excessive toxicity; the higher percentage of bone marrow atrophy reported in control animals than in treated ones; the high mortality among control animals necessitated the use of pooled controls for statistical analysis; the large number of other chemicals tested in the same room and at the same time; and the negative trend for pituitary tumors in treated animals as compared to controls. Despite the experimental shortcomings, the primary reviewer said that the conclusion on the carcinogenicity of Thio-TEPA was still valid. He added that Thio-TEPA would appear to pose a carcinogenic risk to man.

The secondary reviewer agreed with the primary reviewer's critique of the bioassay of Thio-TEPA.


It was moved that the report on the bioassay of Thio-TEPA be accepted as written. The motion was seconded and approved unanimously.

Members present were:

Michael Shimkin (Acting Chairman), University of California
at San Diego
Joseph Highland, Environmental Defense Fund
George Roush, Jr., Monsanto Company
Louise Strong, University of Texas Health Sciences Center
John Weisburger, American Health Foundation

-
- * Subsequent to this review, changes may have been made in the bioassay report either as a result of the review or other reasons. Thus, certain comments and criticisms reflected in the review may no longer be appropriate.

GAYLORD			PRINTED IN U.S.A.



10 Center Drive
Bethesda, MD 20892-1150
301-496-1080

GAYLORD

PRINTED IN U S A

NIH LIBRARY



4 0129 3243

